2 -5 Lang Randot Preschool Stereoacu- 가 가

= =
 가 2 5
 2
 5 96 Randot preschool stereoacu-
 , 84.4% , 2
 96 87% . Randot preschool
 47.4%, 3 stereoacu-
 135 arc sec, 4 48 arc sec, 5 50 arc sec . Randot preschool stereoacu-
 ity sensitivity specificity 100%, 94.2%
 100%, 92.4% . 가 3
 가 가 , Randot
 preschool stereoacu- 3 5
 가 (40:1385 ~ 1390, 1999).

= Abstract =

Assessment of Stereoacu- ity with the Randot Preschool Stereoacu- ity and Lang Test in the 2- to 5-Year Age Range

Se Youp Lee, M.D., Sang Hwan Bae, M.D.

< : 1998 11 18 , : 1999 2 10 >

Address reprint requests to Se Youp Lee, M.D.
 Department of Ophthalmology, College of Medicine, Keimyung University
 #194 Dongsan-dong, Chung-gu, Taegu, 700-310, Korea
 Tel : 82-53-250-7720, 7703, Fax : 82-53-250-7705

* 1998 4 80 () .

(1385)217

The evaluation of stereoacuity provides relatively accurate assessment of binocular function. However, it is difficult to perform and also the reliability rate is low in the 2- to 5-year-old age range. We investigated the overall success rate, the success rate according to age, and efficacy with the Randot preschool stereoacuity and Lang test in the 2-to 5-year-old children of 96 patients. The overall success rate of the two stereotests were 84.4%, 47.4% in the 2-year-old children and more than 87% in children over 3-year-old. When children with normal binocular vision were examined with the Randot preschool stereoacuity test, mean stereoacuity was 333 arc sec in the 2-year-age group, 135 arc sec in the 3-year-age group, 48 arc sec in the 4-year-age group, and 50 arc sec in the 5-year-age group. Sensitivity and specificity of Randot preschool stereoacuity test were 100%, 94.2%, respectively. Sensitivity and specificity of Lang test were 100%, and 92.4% respectively. As a result, Randot preschool stereoacuity and Lang stereotests might be considered easy available and valuable test for evaluating binocular function in the 3- to 5-year-old preschool children. Moreover, the Randot preschool stereoacuity provided a relatively accurate measurement of stereoacuity in the 3- to 5-year-old age range(J Korean Ophthalmol Soc 40:1385 ~ 1390, 1999).

Key Words : Lang test, Preschool, Randot preschool stereoacuity test, Stereoacuity

가 , , , stereothreshold 가
 2 5
 . , Randot preschool stereoacuity
 Frisby , 가 가
 1-11) Frisby 3
 stereothreshold sensitivity specificity .
 가 가 가 4),
 가 2).
 1998 4 10
 . 가 96
 500 arc sec 250 arc sec 35 2 5 가 61 , 가
 가 4.5 , , 가
 3-4
 8) .
 Randot preschool stereoacuity 60
 8) , ,
 가

Table 1. Patient Diagnosis

Diagnosis	No.*
Intermittent exotropia	35
Alternating exotropia	6
Nonaccommodative esotropia	7
Partially accommodative esotropia	12
Congenital esotropia	3
Superior oblique palsy	2
Esophoria	1
Amblyopia anisometric	2
strabismic	6
Normal	22
Total	96

*No. : Number of patients

Table 2. Disparities of Randot Preschool Stereoacuity Test

Book Number	Test Position	Disparity (Second of Arc)
1	Top	200
	Bottom	100
2	Top	60
	Bottom	40
3	Top	800
	Bottom	400

Fig. 1. The Randot Preschool Stereoacuity Test.

0.6 , 가
 2 가
 , 가 가
 , 36 (Table 1).
 2 19
 10 , 9 , 3 32
 , 23 ,
 9 , 4 24 ,
 15 , 9
 , 5 21 , 가
 12 , 9 . 가
 Randot preschool stereoacuity (Stereo Optical Company, Chicago, U.S.A.) 3

가 . 1
 200 arc sec 100 arc sec,
 2 60 arc sec 40 arc sec, 3
 800 arc sec 400 arc sec
 (Fig. 1, Table 2).
 1 1 2
 3 . 1
 2 60 arc sec 가 , 1
 sec, 400 arc sec 3 800 arc
 2 ,
 4
 , 2 , 3 가
 (Fig. 1).
 30cm
 가 가 4 가 가
 (motion parallax)
 reflection . 가
 , 3 2
 가 가
 , 가
 가

가 가
 , 가
 . Sensitivity 가
 가 specificity 가
 가 . Randot

preschool stereoacuity
 200 arc sec⁸⁾.
 9.5 × 14.5cm 3가
 3가 가 600 arc sec,
 1200 arc sec, 550 arc sec .
 cm , ,
 가 .

Randot preschool stereoacuity
 2 5 96
 84.4%(81) .
 2 47.4%(9/19), 3 87.5%(28/
 32), 4 100%(24/24), 5 95.2%
 (20/21) .
 84.4%(81)
 2 47.4%(9/19), 3 90.6%(29/32), 4
 100%(24/24), 5 90.5%(19/21)
 (Table 3). Randot preschool stereoacuity
 2 333 arc sec, 3 135 arc sec,
 4 48 arc sec, 5 50 arc sec
 (Table 4). Randot preschool stereoacuity
 sensitivity 100%(29/
 29), 2 100%(3/3), 3 100%
 (8/8), 4 100%(9/9), 5 100%(9/9)
 . Specitivity 94.2%(49/52) , 가

Table 3. Age Group and Success Rates for Stereopsis Assessment with the Randot Preschool Stereoacuity Test and Lang Test

Age(years)	N*	Randot preschool stereoacuity test	Lang test
2.0 to 2.9	19	47.4%(9)	47.4%(9)
3.0 to 3.9	32	87.5%(28)	90.6%(29)
4.0 to 4.9	24	100%(24)	100%(24)
5.0 to 5.9	21	95.2%(20)	90.5%(19)
	96	84.4%(81)	84.4%(81)

*N,() : Number of patients

Table 4. Mean Stereoacuity in Each Age Group of Normal Binocular Vision with the Randot Preschool Stereoacuity Test

Age(years)	N*	Stereoacuity
2.0 to 2.9	6	333
3.0 to 3.9	9	135
4.0 to 4.9	5	48
5.0 to 5.9	2	50
	22	141

*N,() : number of patients

2 83.3%(5/6), 3 90%(18/20), 4
 100%(15/15), 5 100%(11/11) .
 sensitivity 100%
 (27/27) , 2 100%(2/2), 3
 100%(8/8), 4 100% (9/9), 5 100%
 (8/8) . Specitivity 92.4%
 (49/53), 2 83.3%(5/6), 3
 85.7%(18/21), 4 100%(15/15), 5
 100%(11/11) .
 sensitivity specificity

가 6
 7,8,12,13) .

2 4 specificity가 2-3 4-5
 가 가 가 1 , 3 2 가
 가 .
 가 .
 Randot preschool stereoacuity

Randot preschool stereoacuity Randot preschool stereoacuity 가

5

Randot preschool stereoacuity

Randot preschool stereoacuity

가 가⁸⁾
 가³⁾ (pass) (fail)

^{3,8,14)}

Randot preschool stereoacuity

Birch Randot preschool stereoacuity (circle)

2 333 arc sec, 3
 135 arc sec, 4 48 arc sec
 Birch 2 100 200 arc sec, 3
 60 arc sec, 4 40 arc sec

2 5
 , Randot preschool stereoacuity
 86.4% 95.1%

2,3,4 Birch
 가⁸⁾ 2

37.8%,
 48.2%

가 Birch
 1 가 800 arc sec

⁸⁾
 Randot preschool stereoacuity

sensi-
 tivity Birch Randot preschool
 stereoacuity 가

Randot preschool stereoacuity
 84.4% Birch
 2 47.4%

sensitivity가⁸⁾
 sensitivity가

, 3 87.5%
 47.4% 2

Randot preschool stereoacuity
 100%

2 47.4% 2

Randot preschool stereoacuity
 3 5

Randot preschool stereoacuity

2

REFERENCES

- 1) Lang J : A new stereotest. J Pediatr Ophthalmol Strabismus 20:72-74, 1983.
- 2) Simons K : A comparison of the Frisby, Random-Dot E, TNO and Randot circle stereotest in screening and office use. Arch Ophthalmol 99:446-452, 1981.
- 3) Schanel-Klitsch E, Ciner EB, Scheiman M : Stereopsis assessment in normal 1.5 to 3 year old children : A comparison of three random dot tests: Operant preferential looking; Random dot E; and The Lang stereotest I. Binocular vision and strabismus quartly 11: 125-130, 1996.
- 4) Broadbent H, Westall CA : An evaluation of techniques for measuring stereopsis in infants and young children. Ophthalmic Physiol Opt 10:3-7, 1990.
- 5) Frisby JH, Davis H, McMorrow K : An improved training procedure as a precursor to testing young children with the Frisby stereotest. Eye 10:286-290, 1996.
- 6) Birth EE, Gwiazda J, Held R : Stereoacuity development for crossed and uncrossed disparities in human infants. Vis Res 22:507-513, 1982.
- 7) Manny RE, Martinez AT, Fern KD : Testing stereopsis in the preschool child: Is it clinically useful ? J Pediatr Ophthalmol Strabismus 28:223-231, 1991.
- 8) Birch B, Williams C, Hunter J, Lapa MC : Random dot stereoacuity of preschool children J Pediatr Ophthalmol Strabismus 34:217-222, 1997.
- 9) Tyler CW : Sensory processing of binocular disparity In: Schor CM, Cuiffreda KJ, eds. Vergence Eye Movement: Basic and Clinical Aspects. Boston, Mass: Butterworths, 1983, pp. 199-295.
- 10) Lang, J. : New stereo tests. Arch Ophthalmol 31:651-660, 1990.
- 11) Frisby JH, Davis H, McMorrow K : An improved training procedure as a precursor to testing young children with the Frisby stereotest. Eye 10:286-290, 1996.
- 12) Simmermann JS : Absolute threshold of stereopsis using the Frisby test. J Am Optom Assoc 55:50-53, 1984.
- 13) Frisby JP, Davis H, McMorrow K : An improved training procedure as a precursor to testing young children with the Frisby stereotest. Eye 10:286-290, 1996.
- 14) Simon K, Avery KE, Novak A : Small-target random dot stereogram and binocular suppression testing for preschool vision screening. J Pediatr Ophthalmol Strabismus 33:104-113, 1996.