


# Change in management and outcome of mechanical ventilation in Korea: a prospective observational study

Jae Kyeom Sim<sup>1,\*</sup>, Sang-Min Lee<sup>2,\*</sup>, Hyung Koo Kang<sup>3</sup>, Kyung Chan Kim<sup>4</sup>, Young Sam Kim<sup>5</sup>, Yun Seong Kim<sup>6</sup>, Won-Yeon Lee<sup>7</sup>, Sunghoon Park<sup>8</sup>, So Young Park<sup>9</sup>, Ju-Hee Park<sup>10</sup>, Yun Su Sim<sup>11</sup>, Kwangha Lee<sup>12</sup>, Yeon Joo Lee<sup>13</sup>, Jin Hwa Lee<sup>14</sup>, Heung Bum Lee<sup>15</sup>, Chae-Man Lim<sup>16</sup>, Won-Il Choi<sup>17</sup>, Ji Young Hong<sup>18</sup>, Won Jun Song<sup>1,19</sup>, and Gee Young Suh<sup>1,19</sup>

<sup>1</sup>Department of Critical Care Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul; <sup>2</sup>Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, Seoul National University Hospital, Seoul; <sup>3</sup>Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, Inje University Ilsan Paik Hospital, Goyang; <sup>4</sup>Department of Internal Medicine, Daegu Catholic University Medical Center, Daegu; <sup>5</sup>Division of Pulmonology, Department of Internal Medicine, Institute of Chest Disease, Severance Hospital, Yonsei University College of Medicine, Seoul; <sup>6</sup>Division of Pulmonary, Allergy and Critical Care Medicine, Department of Internal Medicine, Pusan National University Yangsan Hospital, Yangsan; <sup>7</sup>Division of Pulmonary, Allergy and Critical Care Medicine, Department of Internal Medicine, Wonju Severance Christian Hospital, Yonsei University Wonju College of Medicine, Wonju; <sup>8</sup>Division of Pulmonary, Allergy and Critical Care Medicine, Department of Internal Medicine, Hallym University Sacred Heart Hospital, Anyang; <sup>9</sup>Department of Pulmonary and Critical Care Medicine, Department of Medicine, Kyung Hee University Medical Center, Seoul; <sup>10</sup>Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, Seoul Metropolitan Government Seoul National University Boramae Medical Center, Seoul; <sup>11</sup>Division of Pulmonary, Allergy and Critical Care Medicine, Department of Internal Medicine, Hallym University Kangnam Sacred Heart Hospital, Seoul; <sup>12</sup>Division of Pulmonary, Allergy and Critical Care Medicine, Department of Internal Medicine, Pusan National University School of Medicine, Busan; <sup>13</sup>Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, Seoul National University Bundang Hospital, Seongnam; <sup>14</sup>Division of Pulmonary and Critical Care Medicine, Department of Medicine, Ewha Womans University School of Medicine, Seoul; <sup>15</sup>Division of Allergy and Respiratory Diseases, Department of Internal Medicine, Jeonbuk National University Hospital, Jeonju; <sup>16</sup>Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, Asan Medical Center, University of Ulsan College of Medicine, Seoul; <sup>17</sup>Department of Internal Medicine, Keimyung University Dongsan Hospital, Daegu; <sup>18</sup>Division of Pulmonary and Critical Care Medicine, Department of Medicine, Hallym University Chuncheon Sacred Heart Hospital, Chuncheon; <sup>19</sup>Division of Pulmonary and Critical Care Medicine, Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea

## Change in management and outcome of mechanical ventilation in Korea: a prospective observational study



Received : June 9, 2020  
Revised : July 28, 2020  
Accepted : August 4, 2020

Correspondence to Gee Young Suh, M.D.

Department of Pulmonary and Critical Care Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, 81 Irwon-ro, Gangnam-gu, Seoul 06351, Korea  
Tel: +82-2-3410-3429, Fax: +82-2-3410-6956, E-mail: suhgy@skku.edu  
https://orcid.org/0000-0001-5473-1712

\*These authors contributed equally to this work.

**Background/Aims:** There are few studies describing contemporary status of mechanical ventilation in Korea. We investigated changes in management and outcome of mechanical ventilation in Korea.

**Methods:** International, prospective observational cohort studies have been conducted every 6 years since 1998. Korean intensive care units (ICUs) participated in 2010 and 2016 cohorts. We compared 2016 and 2010 Korean data.

**Results:** Two hundred and twenty-six patients from 18 ICUs and 275 patients from 12 ICUs enrolled in 2016 and 2010, respectively. In 2016 compared to 2010, use of non-invasive ventilation outside ICU increased (10.2% vs. 2.5%,  $p = 0.001$ ). Pressure-control ventilation was the most common mode in both groups. Initial tidal volume (7.1 mL/kg vs. 7.4 mL/kg,  $p = 0.372$ ) and positive end-expiratory pressure (6 cmH<sub>2</sub>O vs. 6 cmH<sub>2</sub>O,  $p = 0.141$ ) were similar, but peak pressure (22 cmH<sub>2</sub>O vs. 24 cmH<sub>2</sub>O,  $p = 0.011$ ) was lower in 2016. More patients received sedatives (70.7% vs. 57.0%,  $p = 0.002$ ) and analgesics (86.5% vs. 51.1%,  $p < 0.001$ ) in 2016. The awakening (48.4% vs. 31.0%,  $p = 0.002$ ) was more frequently attempted in 2016. The accidental extubation rate decreased to one tenth of what it was in 2010 (1.1% vs. 10.2%,  $p < 0.001$ ). The ICU mortality did not change (31.4% vs. 35.6%,  $p = 0.343$ ) but ICU length of stay showed a decreasing trend (9 days vs. 10 days,  $p = 0.054$ ) in 2016.

**Conclusions:** There were temporal changes in care of patients on mechanical ventilation including better control of pain and agitation, and active attempt of awakening.

**Keywords:** Respiration, artificial; Pain; Agitation; Ventilator weaning

## INTRODUCTION

Mechanical ventilation is one of the most important treatment interventions provided in the intensive care unit (ICU) with about one third of critically ill patients receiving mechanical ventilation for respiratory failure of various causes [1]. Although mechanical ventilation is needed to help patients overcome the acute stages of critical illness, its application is not without risk. There are guidelines based on randomized clinical trials on how to provide mechanical ventilation to patients safely minimizing complications and to improve survival [2]. However, perhaps it is more important to assess whether such strategies are implemented properly in daily practice, thereby providing guidance on how to improve management of these patients.

Esteban et al. [1,3,4] have conducted international cohort studies every 6 years since 1998 to describe the general characteristics, treatment-related variables, and clinical outcomes of patients who have received mechanical ventilation and to evaluate the evolution of management according to newly published evidence. Twelve Korean ICUs participated in the third international cohort for the first time in 2010 [5], and 18 Korean ICUs participated in the fourth cohort in 2016. The aim of this study was to identify meaningful changes in management and outcome of patients undergoing mechanical ventilation in Korea from two cohorts of

2010 and 2016 studies.

## METHODS

### Design and population

Similar to previous studies [1,3,4], the fourth study was a prospective, and multicenter, non-interventional single-cohort study that maintained similar methodologies (<https://clinicaltrials.gov/ct2/show/record/NCT02731898>). Adult patients who were admitted to ICUs between July 1 and July 31, 2016, and required invasive mechanical ventilation longer than 12 hours or non-invasive ventilation (NIV) for more than 1 hour were included. We extracted and analyzed the data of patients from 18 participating Korean ICUs from the fourth cohort (2016 cohort) and compared them with data of Korean ICUs from the third international study (2010 cohort).

### Data collection

Only the investigator and research coordinators at each site were aware of the purpose and the timing of the study to minimize any practice modifications in response to observation. Baseline demographics were recorded at the beginning of mechanical ventilation. Ventilator and clinical parameters, co-adjutant therapies, and complications were

recorded daily for the duration of mechanical ventilation, or until the 28th day. Variables relevant to ventilator liberation and mortality were also collected.

There were changes in the definitions and terminologies of some complications. The American-European Consensus Conference definition was used in the 2010 cohort to define acute respiratory distress syndrome (ARDS) [6], whereas the Berlin definition was applied to the 2016 cohort [7]. However, only moderate to severe patients, based on the newer definition, were considered to have ARDS in order to maintain consistency. Ventilator-associated pneumonia was investigated in the 2016 cohort instead of ICU-acquired pneumonia, but they were conceptually identical and based on the modified criteria of the Centers for Disease Control and Prevention [8]. The terminology for neuromuscular weakness was changed from critical illness polyneuropathy and myopathy in the 2010 cohort to ICU-acquired weakness (ICUAW) in the 2016 cohort [9]. As with the previous study by Jeong et al. [5], organ failure was defined as having a Sequential Organ Failure Assessment (SOFA) score of  $\geq 3$  for each organ [10]. However, the definition of renal failure was altered in our study. We used serum creatinine or daily urine output corresponding to SOFA scores  $\geq 3$  rather than serum creatinine or the need for renal replacement therapy used in previous studies [5]. The definition of sepsis remained unchanged [11].

In this study, 28-day mortality was calculated from the first day of mechanical ventilation. Ventilator-free days (VFDs) were defined as days alive without invasive mechanical ventilation within the 28 days. The study protocol was approved by the Institutional Review Board (IRB) of each hospital (Samsung Medical Center, IRB No. 2016-04-142; Seoul National University Hospital, IRB No. H-1605-109-763), and the need for consent was waived due to the non-interventional nature of the protocol.

### Statistical analysis

Continuous variables were expressed as medians with interquartile ranges, and the categorical variables were expressed as numbers and percentages. The Mann-Whitney *U* test and Fisher's exact test were used to analyze continuous and categorical variables, respectively. All tests were two sided, and *p* value less than 0.05 was considered statistically significant. Data were analyzed using SPSS version 25.0 (IBM, Armonk, NY, USA).

## RESULTS

### Baseline characteristics

A total of 226 patients from 18 Korean ICUs were enrolled in the 2016 cohort, and their data was compared with that of 275 patients from 12 ICUs from the 2010 cohort. Table 1 shows the baseline characteristics of patients in the two cohorts. Age, body mass index, and Simplified Acute Physiology Score II (SAPS II) score on admission were similar in the two cohorts. More patients were male in the 2016 cohort than in the 2010 cohort, but the difference was not significant (63.7% vs. 58.2%,  $p = 0.232$ ). Pneumonia (25.1%) and sepsis (14.7%) were the most common conditions leading to mechanical ventilation in both cohorts, and followed by aspiration (11.0%) in the 2016 cohort (Table 1).

### Management of mechanical ventilation

Table 2 shows variables related to the management of mechanically ventilated patients. The use of NIV before coming to the ICU increased to 23 patients (10.2%) in the 2016 cohort from seven patients (2.5%) in the 2010 cohort ( $p < 0.001$ ). But, only three patients (1.3%) were treated with NIV as their initial ventilator strategy including one patient who was started on NIV before coming to the ICU in the 2016 cohort while in the 2010 cohort, seven (2.5%) patients including two who started NIV outside the ICU received NIV as their initial strategy ( $p = 0.523$ ). Underlying disease for the three patients in the 2016 cohort included one congestive heart failure (CHF) and one chronic obstructive pulmonary disease (COPD) patients. In the 2010 cohort, two had COPD, two had chronic respiratory disease, and two had either pneumonia or aspiration. In the 2016 cohort, NIV failure was not seen while in the 2010 cohort four out of seven patients eventually needed intubation and invasive mechanical ventilation. Prognosis for patients who were initially treated with NIV was good with every patient alive at ICU discharge except for one patient who needed invasive mechanical ventilation in the 2010 cohort.

For both cohorts, pressure-control ventilation (PCV) was the most frequently applied mode, followed by pressure-support ventilation (PSV), and volume-cycled assist-control ventilation (ACMV). The proportion of patients with PCV was particularly high in the early period, and gradual and reciprocal switch in the proportion of patients with PCV and PSV was found (Fig. 1). Although ACMV remained the third most common mode in the 2016 cohort, its use

**Table 1. Baseline characteristics**

Characteristic	2016 (n = 226)	2010 (n = 275)	p value
Age, yr	68.0 (58.0–77.0)	69.0 (54.0–75.0)	0.250
Male sex <sup>a</sup>	144/226 (63.7)	160/275 (58.2)	0.232
BMI, kg/m <sup>2</sup>	22.0 (20.0–25.0)	22.0 (19.0–24.0)	0.343
SAPS II	51.5 (40.0–62.0)	50.0 (38.0–64.0)	0.610
Previous tracheostomy <sup>a</sup>	8/226 (3.5)	6/275 (2.2)	0.420
NIV at home <sup>a</sup>	2/226 (0.9)	3/275 (1.1)	1.000
Reason for mechanical ventilation <sup>a</sup>			
Acute on chronic respiratory failure			
COPD	6/191 (3.1)	23/275 (8.4)	0.030
Asthma	3/191 (1.6)	3/275 (1.1)	0.693
Other chronic respiratory diseases	9/191 (4.7)	19/275 (6.9)	0.429
Acute respiratory failure			
ARDS	16/191 (8.4)	27/275 (9.8)	0.629
Postoperative	1/191 (0.5)	12/275 (4.4)	0.018
Congestive heart failure	18/191 (9.4)	16/275 (5.8)	0.151
Aspiration	21/191 (11.0)	14/275 (5.1)	0.020
Pneumonia	48/191 (25.1)	63/275 (22.9)	0.582
Sepsis	28/191 (14.7)	28/275 (10.2)	0.150
Trauma	1/191 (0.5)	2/275 (0.7)	1.000
Cardiac arrest	10/191 (5.2)	15/274 (5.5)	1.000
Others <sup>b</sup>	7/191 (3.7)	29/274 (10.5)	0.007
Coma <sup>c</sup>			
Metabolic	10/20 (50.0)	8/22 (36.4)	
Intoxication	4/20 (20.0)	3/22 (13.6)	
Hemorrhagic stroke	2/20 (10.0)	6/22 (27.3)	
Ischemic stroke	1/20 (5.0)	5/22 (22.7)	
Brain trauma	2/20 (10.0)	0/22 (0)	
Others	1/20 (5.0)	0/22 (0)	
Neuromuscular disease	3/191 (1.6)	1/275 (0.4)	0.310

Values are presented as median (interquartile range) or number (%).

BMI, body mass index; SAPS II, Simplified Acute Physiology Score II; NIV, non-invasive ventilation; COPD, chronic obstructive pulmonary disease; ARDS, acute respiratory distress syndrome.

<sup>a</sup>Because of missing data, available number of patients were presented as denominator or in bracket.

<sup>b</sup>Other causes of acute respiratory failure include pulmonary embolism, upper airway obstruction, gastrointestinal bleeding, pneumothorax, hemoptysis and vasculitis, and pulmonary embolism, upper airway obstruction and gastrointestinal bleeding, pulmonary hemorrhage and shock in 2016 and in 2010, respectively.

<sup>c</sup>The percentage of coma subtype was obtained based on the total number of coma.

had decreased significantly compared to 2010 cohort (Table 2) in contrast to the increased use of PCV and PSV which occupied 84% of the total period of invasive mechanical ventilation in 2016. Synchronized intermittent mandatory

ventilation with pressure-support was seldom employed in the 2016 cohort.

Median tidal volume per predicted body weight (TV/PBW) on the first day of mechanical ventilation was 7.1 mL/kg for

**Table 2. Management and co-adjuvant therapy of mechanical ventilation**

Variable	2016 (n = 226)	2010 (n = 275)	p value
NIV as initial ventilator at ICU <sup>a</sup>	3/226 (1.3)	7/275 (2.5)	0.523
NIV success at ICU	3/3 (100)	3/7 (42.9)	0.200
NIV before ICU <sup>a</sup>	23/226 (10.2)	7/275 (2.5)	< 0.001
NIV as initial ventilator at ICU	1/23 (4.3)	2/7 (28.6)	0.128
Ventilator mode <sup>b</sup> (days of use per 1,000 days)			
ACMV	86	205	< 0.001
PSV	300	215	< 0.001
SIMV	19	12	0.058
SIMV + PS	2	143	< 0.001
PCV	540	376	< 0.001
APRV	22	4	< 0.001
PRVC	14	33	< 0.001
High frequency	1	0	0.433
CPAP	15	0	< 0.001
ASV	1	9	< 0.001
Others	0	1	0.263
Ventilator setting at the first day <sup>a,b</sup>			
TV/PBW, mL/kg	7.1 (6.3–8.5) [210]	7.4 (6.2–8.8) [259]	0.372
Peak pressure, cmH <sub>2</sub> O	22.0 (17.8–26.0) [190]	24.0 (19.0–28.0) [249]	0.011
PEEP, cmH <sub>2</sub> O	6.0 (5.0–8.0) [214]	6.0 (5.0–8.0) [265]	0.141
Sedatives <sup>a,b</sup>			
Midazolam	62/222 (27.9)	129/272 (47.4)	< 0.001
Lorazepam	20/222 (9.0)	31/272 (11.4)	0.458
Propofol	37/222 (16.7)	26/272 (9.6)	0.021
Dexmedetomidine	75/222 (33.8)	0/272 (0)	< 0.001
Analgesics <sup>a,b</sup>			
Morphine	21/222 (9.5)	53/272 (19.5)	0.002
Fentanyl	46/222 (20.7)	108/272 (39.7)	< 0.001
Remifentanyl	157/222 (70.7)	30/272 (11.0)	< 0.001
Awakening trial <sup>c</sup>	76/157 (48.4)	48/155 (31.0)	0.002
Neuromuscular blocker <sup>a,b</sup>	45/222 (20.3)	71/272 (26.1)	0.136
Prone position <sup>a,b</sup>	5/222 (2.3)	4/272 (1.5)	0.738
Steroid <sup>a,b</sup>	58/222 (26.1)	92/272 (33.8)	0.077
Intensive insulin therapy <sup>a,b</sup>	35/222 (15.8)	77/272 (28.3)	0.001


Values are presented as number (%) or median (interquartile range).

NIV, non-invasive ventilation; ICU, intensive care unit; ACMV, volume-cycled assist-control ventilation; PSV, pressure-support ventilation; SIMV, synchronized intermittent mandatory ventilation; PS, pressure-support; PCV, pressure-control ventilation; APRV, airway pressure release ventilation; PRVC, pressure regulated volume control; CPAP, continuous positive airway pressure; ASV, adaptive support ventilation; TV, tidal volume; PBW, predicted body weight; PEEP, positive end-expiratory pressure.

<sup>a</sup>Because of missing data, available number of patients were presented as denominator or in bracket.

<sup>b</sup>Ventilator parameters and co-adjuvant therapies were investigated among patients supported by invasive mechanical ventilation. 223 and 272 patients were supported by invasive mechanical ventilation in 2016 and 2010, respectively.

<sup>c</sup>The awakening trial was conducted among patients who received sedatives.



**Figure 1.** Ventilator mode during the course of invasive mechanical ventilation. Bar graphs represent the daily number of patients who were supported by invasive mechanical ventilation. Line graph represent the daily proportion of each mode of ventilation. Miscellaneous mode includes airway pressure release ventilation, pressure regulated volume control, continuous positive airway pressure, adaptive support ventilation, and high frequency oscillatory ventilator mode. PCV, pressure-control ventilation; ACMV, volume-cycled assist-control ventilation; PSV, pressure-support ventilation; SIMV, synchronized intermittent mandatory ventilation.


the 2016 cohort which was slightly decreased from 7.4 mL/kg for the 2010 cohort ( $p = 0.141$ ). Peak pressure on the first day was 22.0 cmH<sub>2</sub>O for the 2016 cohort which was significantly reduced from 24 cmH<sub>2</sub>O in the 2010 cohort ( $p = 0.011$ ). The median positive end-expiratory pressure (PEEP) on the first day of mechanical ventilation was similar (6.0 cmH<sub>2</sub>O vs. 6.0 cmH<sub>2</sub>O,  $p = 0.141$ ) for both cohorts.

In the 2016 cohort, the proportion of patients who received sedatives (70.7% vs. 57.0%,  $p = 0.002$ ) and analgesics (86.5% vs. 51.1%,  $p < 0.001$ ) was significantly higher than in the 2010 cohort, and the use of dexmedetomidine (33.8% vs. 0%,  $p < 0.001$ ) and remifentanyl (70.7% vs. 11.0%,  $p < 0.001$ ) was markedly increased. They were the most frequently prescribed sedative and analgesic, respectively. The use of benzodiazepines, particularly midazolam, was decreased to 27.9% in 2016 from 47.4% in 2010 ( $p < 0.001$ ). Although more patients were receiving sedatives, awakening trials were conducted more often (48.4% vs. 31.0%,  $p = 0.002$ ) in the 2016 cohort. For the 2016 cohort,

light sedation (Richmond Agitation-Sedation Scale [RASS] -2 to 1) was maintained in more than 50% of patients who were assessed except for the first 5 days, and agitation (RASS 2 to 4) was present in less than 10% of patients during the overall period of mechanical ventilation (Fig. 2).

### Acute respiratory distress syndrome

Patients, whose main reason for mechanical ventilation was ARDS, were compared separately between the two cohorts in terms of ventilator strategy and co-adjutant therapies (Table 3). The number of patients who were ventilated in PCV mode on the first day of mechanical ventilation was increased significantly (93.3% vs. 55.6%,  $p = 0.015$ ) with no patient being ventilated in ACMV. There were no significant differences in TV/PBW, peak pressure or PEEP in the two cohorts (Table 3) The use of co-adjutant therapies including rescue therapies were also similar between 2010 and 2016 cohorts, except for the increased administration of analgesics (100% vs. 55.6%,  $p = 0.001$ ) in the 2016 cohort.



**Figure 2.** Level of sedation during the course of invasive mechanical ventilation. Bar graphs represent the daily number of patients who were supported by invasive mechanical ventilation and were assessed for sedation and agitation using Richmond Agitation-Sedation Scale (RASS). Deep sedation, light sedation, and agitation are defined by RASS -5 to -3, -2 to 1, and 2 to 4, respectively.

### Liberation from mechanical ventilation

Table 4 shows several aspects of ventilator liberation. More patients attempted weaning trials (71.4% vs. 61.4%,  $p = 0.029$ ) and the use of gradual reduction of ventilatory support methods increased (68.9% vs. 10.8%,  $p < 0.001$ ) in the 2016 cohort. The success rate of planned extubation among patients who attempted weaning was similar, but the rate of accidental extubation decreased to almost one-tenth of the 2010 rate (1.1% vs. 10.2%,  $p < 0.001$ ) in the 2016 cohort. The most common reason for reintubation was an increased work of breathing for both 2010 and 2016 cohorts. NIV was applied for prevention or treatment of post-extubation respiratory failure in 7.5% of patients in 2016, which was similar to the 2010 cohort. High-flow nasal cannula (HFNC) was applied in 32.1% of patients in the 2016 cohort which was not available in 2010. Tracheostomy rates were similar, but the median time to tracheostomy

was reduced (7.0 days vs. 8.5 days,  $p = 0.014$ ), and more patients underwent percutaneous dilatational tracheostomy (PDT) (48.1% vs. 11.9%,  $p = 0.002$ ) in the 2016 cohort.

### Clinical outcomes

Table 5 describes the clinical outcomes and complications of mechanical ventilation in the two cohorts. The ICU mortality rate was 31.4% for the 2016 cohort and 35.6% for the 2010 cohort ( $p = 0.343$ ). ICU mortality was also not significantly different for different initial diagnostic groups in the two cohorts. However, the ICU length of stay (LOS) showed a decreasing trend from a median of 9 days in the 2016 cohort from 10 days in the 2010 cohort ( $p = 0.054$ ). VFD was median 18 days in the 2016 cohort compared to 14 days in the 2010 cohort, although this difference did not reach statistical significance ( $p = 0.534$ ). The rates of patients who suffered complications of sepsis (21.9% vs.

**Table 3. Ventilator strategy and co-adjuvant therapy of acute respiratory distress syndrome**

Variable	2016 (n = 16)	2010 (n = 27)	p value
Ventilator mode at the first day <sup>a</sup>			
ACMV	0/15 (0)	6/27 (22.2)	0.073
PSV	1/15 (6.7)	2/27 (7.4)	1.000
SIMV + PS	0/15 (0)	2/27 (7.4)	0.530
PCV	14/15 (93.3)	15/27 (55.6)	0.015
PRVC	0/15 (0)	2/27 (7.4)	0.530
TV/PBW at the first day, mL/kg <sup>a</sup>			
TV/PBW < 6	4/13 (30.8)	8/26 (30.8)	1.000
6 ≤ TV/PBW < 8	5/13 (38.5)	12/26 (46.2)	0.740
8 ≤ TV/PBW < 10	0/13 (0)	3/26 (11.5)	0.538
TV/PBW > 10	4/13 (30.8)	3/26 (11.5)	0.194
Peak pressure at the first day, cmH <sub>2</sub> O <sup>a</sup>	26.0 (20.8–30.0) [14]	27.5 (21.0–32.3) [26]	0.705
PEEP at the first day, cmH <sub>2</sub> O <sup>a</sup>			
PEEP < 5	2/14 (14.3)	2/27 (7.4)	0.596
5 ≤ PEEP ≤ 10	9/14 (64.3)	16/27 (59.3)	1.000
PEEP > 10	3/14 (21.4)	9/27 (33.3)	0.494
Sedatives	16/16 (100)	24/27 (88.9)	0.282
Analgesics	16/16 (100)	15/27 (55.6)	0.001
Neuromuscular blocker	8/16 (50)	11/27 (40.7)	0.752
Prone position	3/16 (18.8)	2/27 (7.4)	0.344
Inhaled NO	1/16 (6.25)	No data	
ECMO	0/16 (0)	No data	

Values are presented as median (interquartile range) or number (%).

ACMV, volume-cycled assist-control ventilation; PSV, pressure-support ventilation; SIMV, synchronized intermittent mandatory ventilation; PS, pressure-support; PRVC, pressure regulated volume control; TV, tidal volume; PBW, predicted body weight; PEEP, positive end-expiratory pressure; NO, nitric oxide; ECMO, extracorporeal membrane oxygenation.

<sup>a</sup>Because of missing data, available number of patients were presented as denominator or in bracket.

50.6%,  $p < 0.001$ ) was significantly lower in the 2016 cohort. Significantly higher proportion of patients was recognized as having delirium (37.4% vs. 8.0%,  $p < 0.001$ ) and ICUAW (5.0% vs. 0.7%,  $p = 0.004$ ) in the 2016 cohort.

## DISCUSSION

In the current study, we describe the latest status of mechanical ventilation in Korea and observed temporal changes in ventilator operations, the management of pain and agitation, and liberation from ventilators between 2016 and 2010 cohorts participating in international multicenter study.

Similar to 2010, the use of NIV in ICU was still low in 2016. This finding is quite different from situations in other countries where 37% of mechanically ventilated patients used NIV as initial ventilator mode in France [12], and 20% of patients started mechanical ventilation with NIV in the United States [13]. It should be noted that NIV use outside ICU has greatly increased in 2016 cohort compared to 2010 cohort. But surprisingly, only one patient who needed ventilator support due to CHF, continued to receive NIV after the patient was transferred to the ICU. Thus increased use of NIV during pre-ICU period may reflect general shortage of ICU resources rather than increased use of NIV for clinically appropriate indications. This is reflected in the fact that only three patients out of 22 patients who received pre-ICU



**Table 4. Liberation from invasive mechanical ventilation**

Variable	2016 (n = 223)	2010 (n = 272)	p value
Weaning try <sup>a</sup>	135/189 (71.4)	167/272 (61.4)	0.029
Method at first attempt <sup>b</sup>			
GRS	93/135 (68.9)	18/167 (10.8)	< 0.001
SBT	42/135 (31.1)	149/167 (89.2)	< 0.001
Planned extubation <sup>b</sup>	104/135 (77.0)	135/167 (80.8)	0.477
Accidental extubation <sup>a,c</sup>	2/181 (1.1)	27/266 (10.2)	< 0.001
Reintubation <sup>d</sup>	21/106 (19.8)	31/162 (19.1)	1.000
After planned extubation	21/104 (20.2)	24/135 (17.8)	0.739
After accidental extubation	0/2 (0)	7/27 (25.9)	1.000
Reintubation within 2 day	14/21 (66.7)	27/31 (87.1)	0.095
NIV after extubation <sup>d</sup>	8/106 (7.5)	16/162 (9.9)	0.663
HFNC after extubation <sup>d</sup>	34/106 (32.1)	No data	
Tracheostomy <sup>a,c</sup>	27/181 (14.9)	42/266 (15.8)	0.894
Before extubation	18/27 (66.7)	31/42 (73.8)	0.592
After reintubation	9/27 (33.3)	11/42 (26.2)	0.592
Time to tracheostomy, day	7.0 (4.0–15.0)	8.5 (13.0–18.0) <sup>e</sup>	0.014
PDT	13/27 (48.1)	5/42 (11.9)	0.002

Values are presented as median (interquartile range) or number (%). GRS method included pressure-support ventilation (PSV), synchronized intermittent mandatory ventilation (SIMV) and SIMV with pressure-support. SBT method included T-piece, continuous positive airway pressure, and PSV less than 7 cmH<sub>2</sub>O.

GRS, gradual reduction of ventilatory support; SBT, spontaneous breathing trial; NIV, non-invasive ventilation; HFNC, high-flow nasal cannula; PDT, percutaneous dilatational tracheostomy.

<sup>a</sup>Because of missing data, available number of patients were presented as denominator or in bracket.

<sup>b</sup>Patients who attempted weaning trial were included for analysis.

<sup>c</sup>Patients who already had tracheostomy were excluded from the analysis. Six and eight patients already had tracheostomy in 2010 and in 2016, respectively.

<sup>d</sup>Patients who experienced either planned or accidental extubation were included for analysis.

<sup>e</sup>One patient was excluded from analysis due to inappropriate data.

NIV but were started on invasive mechanical ventilation in the ICU had conditions associated with high success rate of NIV such as CHF and neuromuscular disease (data are not shown).

Increased use of pressure modes have been reported previously [1]. However, the preference for pressure modes was stronger in Korea [5], and became more pronounced in the 2016 cohort, especially in ARDS patients in whom pressure modes were exclusively used. This is interesting because there is still no evidence that pressure-targeted ventilation is superior to volume-targeted ventilation [14,15].

One of the most remarkable findings in this study is the temporal changes in the use of analgesics and sedatives. Analgesics were administered in 27% of mechanically ven-

tilated patients in Korea in an unrelated multicenter observational study in 2003 [16]. The prevalence of analgesic use increased to 51.1% in the 2010 cohort [5] and to 86.5% in the 2016 cohort. These changes may reflect that many ICUs now follow the 2013 Pain, Agitation, and Delirium (PAD) guideline, which stressed that appropriate pain control is an important part of patient care [17]. The most frequently used analgesic has changed from morphine to fentanyl [5,16,18] and remifentanyl. Although the 2013 PAD guideline did not recommend a specific opioid, remifentanyl is a reasonable choice as an analgesics in patients on mechanical ventilation because it is easily titratable, dose not accumulate in organ failure, and has short-action time [19]. The proportion of patients who receiving dexmedetomidine and

**Table 5. Clinical outcomes and complications of mechanical ventilation**

Variable	2016 (n = 226)	2010 (n = 275)	p value
ICU mortality	71/226 (31.4)	98/275 (35.6)	0.343
Pneumonia	20/48 (41.7)	20/63 (31.7)	0.322
Sepsis	14/28 (50.0)	12/28 (42.9)	0.789
Aspiration	2/21 (9.5)	4/14 (28.6)	0.191
ARDS	8/16 (50.0)	17/27 (63.0)	0.526
28-Day mortality <sup>a</sup>	68/192 (35.4)	93/273 (34.1)	0.767
Hospital mortality <sup>a</sup>	88/200 (44.0)	112/269 (41.6)	0.637
ICU LOS, day	9.0 (5.0–15.0)	10.0 (5.0–19.0)	0.054
Hospital LOS <sup>a</sup> , day	21.0 (11.0–42.0) [199]	23.0 (12.0–41.0) [269]	0.528
Ventilator-free days	18.00 (0.0–23.0)	14.0 (0.0–23.0)	0.534
Complications <sup>a</sup>			
Sepsis	48/219 (21.9)	132/261 (50.6)	< 0.001
RRT	57/219 (26.0)	62/261 (23.8)	0.596
ARDS	42/219 (19.2)	59/270 (21.9)	0.501
VAP	20/219 (9.1)	20/239 (8.4)	0.869
Barotrauma	5/219 (2.3)	11/261 (4.2)	0.311
Organ failure			
Cardiovascular	140/219 (63.9)	172/261 (65.9)	0.701
Respiratory	175/214 (81.8)	216/268 (80.6)	0.815
Hematologic	73/209 (34.9)	89/248 (35.9)	0.845
Renal	78/216 (36.1)	85/251 (33.9)	0.627
Hepatic	36/209 (17.2)	27/245 (11.0)	0.076
ICU-acquired weakness	11/219 (5.0)	2/275 (0.7)	0.004
Delirium	82/219 (37.4)	22/275 (8.0)	< 0.001

Values are presented as median (interquartile range) or number (%).

ICU, intensive care unit; ARDS, acute respiratory distress syndrome; LOS, length of stay; RRT, renal replacement therapy; VAP, ventilator-associated pneumonia.

<sup>a</sup>Because of missing data, available number of patients were presented as denominator or in bracket.

propofol increased while use of benzodiazepines decreased also reflecting compliance with the 2013 PAD guideline. Also, higher proportion of patients were undertaking awakening trials in 2016 to decrease amount of sedatives.

In the 2016 cohort, proportion of patients using spontaneous breathing trial (SBT) as the method at first attempt was only 31.1%. This was even decreased compared to 2010 cohort which was 89.2%. It is well known that daily screening and SBT can improve outcome such as increasing VFD [20]. In many ICUs around the world, weaning process is protocolized and screening and SBT are carried out by specialized team usually consisting of respiratory therapists.

Since in Korea, there are no respiratory therapists or specialized teams to carry out this task, in many ICUs weaning process is not standardized and thus high percentage of patients weaned through decreased support level as seen with this cohort. This is an area that critical care community as a whole need to improve on.

Since its introduction between 2010 and 2016, HFNC has become widely used in Korean ICU due to several studies favoring its use in acute respiratory failures and post-extubation conditions [21-23]. Until 2010, PDT was not widely adopted in Korea in contrast to international cohorts in whom more than 50% of patients underwent PDTs in 2004 [4]. In

2016, half of all tracheostomies were performed using the percutaneous method.

The changes seen in practice of care in ventilated patients in this study might reflect efforts of the critical society in Korea to bring about change in governmental policy and to educate and cultivate critical care specialist. After years of effort, subspecialty board in critical care was established in 2009 [24]. The government insurance agency, Health Insurance Review and Assessment Service, does periodic assessment of quality of care in critical care which is bound to reimbursement. Also for the first time, hospitals much have full-time intensivists to apply for tertiary-referral center status. Through these efforts, the proportion of ICUs with dedicated intensivists increased from 17.3% in 2009 to 51.1% in 2014 [25].

Despite meaningful changes in the practice of mechanical ventilation, major clinical outcomes such as mortality, LOS, and VFD were not significantly improved. This may be because other confounding factors such as comorbidity or frailty might have influenced outcome. For example, a recent study demonstrated that the Charlson Comorbidity Index has increased over time in Korean ICU patients [26]. Also, the most prominent change between 2010 and 2016 cohorts was widespread implementation of the 2013 PAD guideline. However, the guideline and supporting evidences are not primarily intended to improve survival in ICU but more related to quality of life in the post-ICU period which was not measured. In 2013, the beneficial effects of a prone position were demonstrated in selected ARDS patients [27], and indeed the proportion of ARDS patients who underwent prone position increased in 2016. However, the number of prone positioned patients was too small to affect mortality of the whole study population. In a similar way, management improvement in certain diseases, if any, would not change outcomes of this heterogeneous population. Moreover, TV/PBW, the oldest and strongest determinant of outcome of mechanical ventilation, has been maintained in acceptable range without significant change. It is worth noting the decreasing trend of ICU LOS. This change may result from the improvement of pain and agitation control, and the increase in awakening and weaning trials.

More patients were reported to have ICUAW and delirium in 2016. For the ICUAW, direct comparisons were limited due to varying definitions in 2010 and 2016. The incidence reported in 2016 (5.0%) was much lower than what has been presented in other studies [28], in spite of its rapid rise.

It appears that ICUAWs had been underestimated and the awareness has increased rather than the occurrence. This explanation also fits the increased incidence of delirium. Similar findings were observed in the United States/Canada regions in the fourth international cohort [29].

There are several limitations in the present study. First, we compared the mortality of the overall study population without considering site effects. However, comparing only identical ICUs would lower the statistical power because of the reduced number of patients. All participating institutions were academic teaching hospitals in one country, so site effects may not be large. Second, while the number of participating ICUs in 2016 was larger, the number of enrolled patients was smaller. In addition, missing data were more common with possible concerns regarding the risk of data distortion. However, our result was concordant with the international data [29]. Thirdly, while we evaluated whether a patient received medication, we did not investigate dose or treatment duration.

In conclusion, the current study shows the evolution of management of mechanically ventilated patients in Korea over 6 years. Better control of pain and agitation, and active attempt of awakening was noted but steps to improve weaning process is needed. In addition, the recognition of delirium and ICUAW has improved.

## KEY MESSAGE

1. More patients received sedatives and analgesics. The use of dexmedetomidine and remifentanyl increased. Conversely, the use of midazolam decreased.
2. More patients attempted to awakening and weaning trials.
3. The recognition of delirium and intensive care unit (ICU)-acquired weakness has improved.
4. The ICU mortality did not change but the ICU length of stay showed a tendency to decrease.

## Conflict of interest

No potential conflict of interest relevant to this article was reported.

## Acknowledgments

We appreciate all members of the Korean Study Group on Respiratory Failure (KOSREF).

## REFERENCES

1. Esteban A, Frutos-Vivar F, Muriel A, et al. Evolution of mortality over time in patients receiving mechanical ventilation. *Am J Respir Crit Care Med* 2013;188:220-230.
2. Fan E, Del Sorbo L, Goligher EC, et al. An official American Thoracic Society/European Society of Intensive Care Medicine/Society of Critical Care Medicine Clinical Practice guideline: mechanical ventilation in adult patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2017;195:1253-1263.
3. Esteban A, Ferguson ND, Meade MO, et al. Evolution of mechanical ventilation in response to clinical research. *Am J Respir Crit Care Med* 2008;177:170-177.
4. Esteban A, Anzueto A, Frutos F, et al. Characteristics and outcomes in adult patients receiving mechanical ventilation: a 28-day international study. *JAMA* 2002;287:345-355.
5. Jeong BH, Suh GY, An JY, et al. Clinical demographics and outcomes in mechanically ventilated patients in Korean intensive care units. *J Korean Med Sci* 2014;29:864-870.
6. Bernard GR, Artigas A, Brigham KL, et al. The American-European Consensus Conference on ARDS. Definitions, mechanisms, relevant outcomes, and clinical trial coordination. *Am J Respir Crit Care Med* 1994;149(3 Pt 1):818-824.
7. ARDS Definition Task Force, Ranieri VM, Rubenfeld GD, et al. Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 2012;307:2526-2533.
8. Garner JS, Jarvis WR, Emori TG, Horan TC, Hughes JM. CDC definitions for nosocomial infections, 1988. *Am J Infect Control* 1988;16:128-140.
9. Fan E, Cheek F, Chlan L, et al. An official American Thoracic Society Clinical Practice guideline: the diagnosis of intensive care unit-acquired weakness in adults. *Am J Respir Crit Care Med* 2014;190:1437-1446.
10. Vincent JL, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med* 1996;22:707-710.
11. Bone RC, Balk RA, Cerra FB, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest* 1992;101:1644-1655.
12. Demoule A, Chevret S, Carlucci A, et al. Changing use of non-invasive ventilation in critically ill patients: trends over 15 years in francophone countries. *Intensive Care Med* 2016;42:82-92.
13. Maheshwari V, Paioli D, Rothaar R, Hill NS. Utilization of non-invasive ventilation in acute care hospitals: a regional survey. *Chest* 2006;129:1226-1233.
14. Rittayamai N, Katsios CM, Beloncle F, Friedrich JO, Mancebo J, Brochard L. Pressure-controlled vs volume-controlled ventilation in acute respiratory failure: a physiology-based narrative and systematic review. *Chest* 2015;148:340-355.
15. Chacko B, Peter JV, Tharyan P, John G, Jeyaseelan L. Pressure-controlled versus volume-controlled ventilation for acute respiratory failure due to acute lung injury (ALI) or acute respiratory distress syndrome (ARDS). *Cochrane Database Syst Rev* 2015;1:CD008807.
16. Jang HJ, Ra SW, Oh BJ, Lim CM, Koh Y, Hong SB. Multicenter prospective observational study about the usage patterns of sedatives, analgesics and neuromuscular blocking agents in the patients requiring more than 72 hours mechanical ventilation in intensive care units of Korea. *Korean J Crit Care Med* 2009;24:145-151.
17. Barr J, Fraser GL, Puntillo K, et al. Clinical practice guidelines for the management of pain, agitation, and delirium in adult patients in the intensive care unit. *Crit Care Med* 2013;41:263-306.
18. Kim HH, Choi SC, Ahn JH, Chae MK, Heo J, Min YG. Analysis of trends in usage of analgesics and sedatives in intensive care units of South Korea: a retrospective nationwide population-based study. *Medicine (Baltimore)* 2018;97:e12126.
19. Muellejans B, Lopez A, Cross MH, Bonome C, Morrison L, Kirkham AJ. Remifentanyl versus fentanyl for analgesia based sedation to provide patient comfort in the intensive care unit: a randomized, double-blind controlled trial [ISRCTN43755713]. *Crit Care* 2004;8:R1-R11.
20. Girard TD, Kress JP, Fuchs BD, et al. Efficacy and safety of a paired sedation and ventilator weaning protocol for mechanically ventilated patients in intensive care (Awakening and Breathing Controlled trial): a randomised controlled trial. *Lancet* 2008;371:126-134.
21. Maggiore SM, Idone FA, Vaschetto R, et al. Nasal high-flow versus Venturi mask oxygen therapy after extubation. Effects on oxygenation, comfort, and clinical outcome. *Am J Respir Crit Care Med* 2014;190:282-288.

22. Rittayamai N, Tscheikuna J, Rujiwit P. High-flow nasal cannula versus conventional oxygen therapy after endotracheal extubation: a randomized crossover physiologic study. *Respir Care* 2014;59:485-490.
23. Frat JP, Thille AW, Mercat A, et al. High-flow oxygen through nasal cannula in acute hypoxemic respiratory failure. *N Engl J Med* 2015;372:2185-2196.
24. Lim CM, Kwak SH, Suh GY, Koh Y. Critical care in Korea: present and future. *J Korean Med Sci* 2015;30:1540-1544.
25. The Korean Society of Critical Care Medicine, Committee on White Paper Publication. Daehanjunghwanjauihaghoe baegseo: Je 2 ho 2015 gugnae junghwanjasil hyeonhwangjosa bogoseo [The Korean Society of Critical Care Medicine White Paper: report on the status of intensive care units in Korea No.2 2015]. 2nd ed. Seoul (KR): The Korean Society of Critical Care Medicine, 2016.
26. Heo J, Hong Y, Han SS, et al. Changes in the characteristics and long-term mortality rates of intensive care unit patients from 2003 to 2010: a nationwide population-based cohort study performed in the Republic of Korea. *Acute Crit Care* 2018;33:135-145.
27. Guerin C, Reigner J, Richard JC, et al. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med* 2013;368:2159-2168.
28. Farhan H, Moreno-Duarte I, Latronico N, Zafonte R, Eikermann M. Acquired muscle weakness in the surgical intensive care unit: nosology, epidemiology, diagnosis, and prevention. *Anesthesiology* 2016;124:207-234.
29. Owen GD, Stollings JL, Rakhit S, et al. International analgesia, sedation, and delirium practices: a prospective cohort study. *J Intensive Care* 2019;7:25.