



## *Correction* **Correction:** Kim et al. The Medium Cut-Off Membrane Does Not Lower Protein-Bound Uremic Toxins. *Toxins* 2022, 14, 779

Yang Gyun Kim <sup>1</sup>, Sang Ho Lee <sup>1</sup>, Su Woong Jung <sup>1</sup>, Gun Tae Jung <sup>2</sup>, Hyun Ji Lim <sup>2</sup>, Kwang Pyo Kim <sup>2,3</sup>, Young-Il Jo <sup>4</sup>, KyuBok Jin <sup>5</sup> and Ju Young Moon <sup>1,\*</sup>

- <sup>1</sup> Division of Nephrology, Department of Internal Medicine, Kyung Hee University School of Medicine, Seoul 05278, Republic of Korea
- <sup>2</sup> Department of Biomedical Science and Technology, Kyung Hee Medical Science Research Institute, Kyung Hee University, Seoul 02453, Republic of Korea
- <sup>3</sup> Department of Applied Chemistry, Institute of Natural Science, Global Center for Pharmaceutical Ingredient Materials, Kyung Hee University, Yongin 17104, Republic of Korea
- <sup>4</sup> Division of Nephrology, Department of Internal Medicine, Konkuk University Medical Center, Seoul 05029, Republic of Korea
- <sup>5</sup> Division of Nephrology, Department of Internal Medicine, Keimyung University Dongsang Hospital, Keimyung University Kidney Institute, Daegu 42601, Republic of Korea
- \* Correspondence: jymoon@khu.ac.kr

## Error in Table

In the original publication [1], there was a mistake in Table 2 as published. The information for dialyzers including inner diameter, wall thickness, effective surface area, clearance of inulin, and UF coefficient was incorrect. The corrected Table 2 appears below. The authors state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

Table 2. The characteristics of the dialyzers.

	HF-HD	Post-OL-HDF	MCO-HD
Dialyzer	Fx CorDiax 80	Fx CorDiax 800	Theranova 400
Inner diameter (µm)	185	210	180
Wall thickness (µm)	35	35	35
Membrane polymer	polysulphone-PVP blend	polysulphone-PVP blend	polyarylethersul- phone-PVP blend
Effective surface area (m <sup>2</sup> )	1.8	2.0	1.7
Sieving coefficients of albumin	< 0.001	<0.001	0.008
Clearance of inulin	135	178	183
UF coefficient (mL/h/mmHg)	64	62	48

PVP, polyvinylpyrrolidone; UF, ultrafiltration.

## Reference

 Kim, Y.G.; Lee, S.H.; Jung, S.W.; Jung, G.T.; Lim, H.J.; Kim, K.P.; Jo, Y.-I.; Jin, K.; Moon, J.Y. The Medium Cut-Off Membrane Does Not Lower Protein-Bound Uremic Toxins. *Toxins* 2022, 14, 779. [CrossRef]

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.



Citation: Kim, Y.G.; Lee, S.H.; Jung, S.W.; Jung, G.T.; Lim, H.J.; Kim, K.P.; Jo, Y.-L; Jin, K.; Moon, J.Y. Correction: Kim et al. The Medium Cut-Off Membrane Does Not Lower Protein-Bound Uremic Toxins. *Toxins* 2022, 14, 779. *Toxins* 2023, 15, 64. https://doi.org/10.3390/ toxins15010064

Received: 26 December 2022 Accepted: 9 January 2023 Published: 11 January 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/).