

Minireview

Interaction between epidermal growth factor receptor– and cyclooxygenase 2–mediated pathways and its implications for the chemoprevention of head and neck cancer

Mi Sun Choe,¹ Xin Zhang,¹ Hyung Ju C. Shin,² Dong M. Shin,¹ and Zhuo (Georgia) Chen¹

¹Department of Hematology/Oncology, Winship Cancer Institute, Emory University and ²Quest Diagnostics, Atlanta, Georgia

Abstract

Head and neck squamous cell carcinoma is a well-known model for chemoprevention studies because of its field cancerization effect, its multistep carcinogenesis process, and the easy accessibility of biopsies to target lesions. With new understandings of head and neck carcinogenesis and the development of molecular targeted therapy, chemoprevention trials for head and neck squamous cell carcinoma have been rapidly updated. Cyclooxygenase-2 (COX-2) and epidermal growth factor receptor (EGFR) tyrosine kinase inhibitors are gaining significant attention as potential chemopreventive agents. Both COX-2 and EGFR are involved in head and neck carcinogenesis. Targeting COX-2 and EGFR separately has shown promising antitumor activity. Recently, combinations of COX-2 and EGFR tyrosine kinase inhibitors have been reported to show synergistic/additive effects in preclinical studies. Because COX-2 and EGFR tyrosine kinase inhibitors are toxic as single agents in clinical trials, the combination of COX-2 and EGFR tyrosine kinase inhibitors used at lower doses seems more promising than monotherapy with either as a novel strategy in head and neck cancer chemoprevention. [Mol Cancer Ther 2005;4(9):1448–55]

Introduction

Approximately 40,000 cases of new head and neck cancer (HNC) develop annually, accounting for ~3% of all new cancers in the United States (1), and 90% of these are squamous cell carcinoma [head and neck squamous cell carcinoma (HNSCC)]. Due to its location and anatomic complexity, HNC causes almost inevitable functional and social impairment even before it becomes life threatening. Despite great advances in therapy, the overall survival rate for patients with HNC has not improved significantly, emphasizing the importance of preventive intervention (2–4).

Chemoprevention can be defined as the use of specific agents to suppress, reverse, or prevent carcinogenesis, thus stopping the progression to invasive cancer by modulating the carcinogenic process or by removal (apoptosis) of premalignant cells (5, 6). Recently, the concept of chemoprevention has been substantially incorporated into cancer treatment goals. Progress in clinical trials has shown that the combined preventive approach is more effective than single-agent chemoprevention.

This review will briefly summarize chemopreventive approaches in HNSCC before highlighting a novel and promising chemopreventive modality combining cyclooxygenase-2 (COX-2) and epidermal growth factor receptor (EGFR) tyrosine kinase inhibitors.

Chemoprevention for HNC

HNSCC is an excellent model for the chemopreventive approach in several aspects. First, HNSCC is notorious for its high tendency to develop second primary tumors, one of the main reasons for the typical dismal outcome of this cancer. The lifetime cumulative risk of second primary tumor is >20% and has been reported in up to 47% in patients with previously treated laryngeal cancer (2, 7). This multicentricity is best explained by the fact that the whole mucosa of the upper aerodigestive tract is affected by the same carcinogens, most likely tobacco and alcohol (field cancerization; ref. 8). Second, HNSCC well exemplifies multistep carcinogenesis with stepwise accumulations of genetic alterations (9, 10). Pathologically, the course follows from normal epithelium, hyperplasia, and dysplasia to invasive HNSCC. Multicentricity and multistep carcinogenesis provides the rational background for a chemopreventive

Received 9/20/04; revised 7/13/05; accepted 7/15/05.

Grant support: NIH grant U01 CA101244.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Zhuo (Georgia) Chen, Department of Hematology/Oncology, Winship Cancer Institute, Emory University, Room 3086, 1365-C Clifton Road, Atlanta, GA 30322.
E-mail: georgia_chen@emoryhealthcare.org

Copyright © 2005 American Association for Cancer Research.

doi:10.1158/1535-7163.MCT-04-0251

approach. In addition, HNC exhibits clinically well defined premalignant lesions, such as oral leukoplakia and erythroplakia, to be targeted. These can be easily identified compared with premalignant lesions in other organ sites and provide accessibility for macroscopic and microscopic evaluation or biomarker evaluation of lesions.

HNSCC has been one of the main models for chemopreventive approaches and important issues have been learned from chemopreventive clinical trials. The most extensively studied chemopreventive agents in HNC are retinoids (11). The first randomized clinical trial was conducted by Hong et al. (11) using a high dose of 13-*cis*-retinoic acid (13-cRA) for a short period in patients with oral premalignant lesions, which showed significant clinical and pathologic responses. In spite of the encouraging results, this trial raised two important issues common in chemopreventive clinical trials. Dose-related toxicity was an important obstacle to treating patients on a long-term basis, and the remission induced by short-term treatment did not last long after cessation of treatment. This observation led to a subsequent low-dose maintenance trial. Lippman et al. (5) treated patients with oral premalignant lesions with high-dose 13-cRA for 3 months and switched to either low-dose 13-cRA or β -carotene for maintenance. Although the results showed that using low-dose 13-cRA for maintenance could effectively repress disease progression, the long-term follow-up failed to show a difference in the cancer development rate in both groups (11). In terms of prevention of second primary tumor, the first phase III clinical trial using 13-cRA showed a statistically significant suppression of second primary tumor, although there was no significant difference in survival. Subsequent randomized clinical trials using retinoids have failed to show a difference in the development of second primary tumor or survival (11).

Obtaining long-lasting efficacy with low toxicity has been an important issue, provoking combinations of retinoids with other agents or searches for new chemopreventive agents. Other chemopreventive agents that have shown preventive effects include selenium and vitamin E. Recently, Papadimitrakopoulou et al. (12) reported that a combination of 13-cRA, vitamin E, and IFN- α restored advanced laryngeal premalignant lesions. Shin et al. (13, 14) showed that the same combination suppressed the development of second primary tumor and/or recurrence, and achieved an excellent survival rate in stage III/IV HNSCC, making this a promising combination chemopreventive approach. With a better understanding of relevant molecular changes in each carcinogenic step of HNC and increased availability of specific molecular targeting agents, chemoprevention in HNC has evolved to a new era. New agents in this context include COX inhibitors, EGFR tyrosine kinase inhibitors, farnesyl transferase inhibitors, and others (11).

Rationale for Blocking COX-2 and EGFR Pathways in Chemoprevention of HNC

Selection of chemopreventive agents requires consideration of several criteria. Biological efficacy impeding the

carcinogenic process, selectivity for transformed tissue, minimal toxicity, and the ability to obtain good patient compliance are essential. As new potential agents, COX-2 inhibitor and EGFR inhibitor show promise for chemoprevention of HNC. Both the COX-2 and EGFR signaling pathways play major roles in head and neck carcinogenesis (15, 16). Both COX-2 and EGFR are overexpressed in premalignant and malignant tissues of the head and neck compared with normal tissue (17–20). Blocking these pathways has already shown promising antitumor activity. In addition, both the COX-2 and EGFR inhibitors used in clinics are orally bioavailable and their side effects are well tolerated at clinically relevant dosages (21–23).

COX-2 Pathway in Carcinogenesis

COX-1 and COX-2 catalyze prostanoid synthesis from arachidonic acid. COX-1 is constitutively expressed in nearly all normal tissues and has a beneficial housekeeping role. COX-2 is undetectable (or at low levels) in most normal tissues but is rapidly induced in response to inflammatory or mitogenic stimuli, including cytokines, growth factors, and tumor promoters (24, 25).

COX-2 has been implicated in carcinogenesis ever since the discovery that intake of nonsteroidal anti-inflammatory drugs that inhibit COX activity also decrease the relative risk for development of colorectal cancer (26). There is also direct evidence for the carcinogenic role of COX-2. Oshima et al. (27) showed that selective genetic elimination of COX-2 protected APC tumor suppressor gene-deleted mice from developing intestinal tumors. In humans, COX-2 overexpression has been found in premalignant and malignant lesions in several tumors (17, 18, 25; for review, see 28). In HNC, there is a stepwise increase in COX-2 expression through the normal, hyperplastic, dysplastic, and invasive carcinoma stages, suggesting its carcinogenic role in HNC (17, 18, 29).

COX-2 overexpression contributes to many aspects of carcinogenesis, such as inhibition of apoptosis, promotion of cell proliferation, induction of angiogenesis, and increasing invasiveness, mainly through increasing the amount of prostaglandins, including prostaglandin E₂ (PGE₂), PGF_{2 α} , PGD₂, TXA₂, PGI₂, and PGJ₂ (28, 30). Prostaglandins exert their effect mainly by binding to G-protein-coupled receptors, including EP-1, EP-2, EP-3, and EP-4, on cell surfaces (for review, see ref. 31). Each is specifically recognized by certain prostaglandins and activates signaling transduction from extracellular signal regulated kinase (32), phosphatidylinositol 3-kinase/AKT, and cyclic AMP/protein kinase A (32–35). Alternatively, cyclopentenone prostaglandins, such as PGJ₂, can bind to the nuclear transcription factor peroxisome proliferator-activated receptor, which regulates expression of several genes involved in cell proliferation (36).

EGFR Signaling Pathway in Cancer Progression

EGFR, a surface receptor with intrinsic tyrosine kinase activity, is one of several known pivotal intermediates in many epithelial malignancies (37). It belongs to the erbB growth factor receptor family. The ligands binding to the

extracellular domain of EGFR induce homodimerization of EGFR or heterodimerization with other members of the erbB growth factor receptor family, activating the intrinsic tyrosine kinase and its downstream signaling molecules (38). Potential EGFR downstream signaling pathways include Ras/mitogen-activated protein kinase, phosphatidylinositol 3-kinase, phospholipase C γ , the Src kinase family, Janus kinase, signal transducers, activators of transcription, and others (39). Activation of EGFR is involved in pertinent pleiotropic cellular processes, such as cell proliferation, apoptosis, differentiation, angiogenesis, and motility (37–43).

Overexpression of EGFR has been frequently reported in human malignant neoplasms (44). In HNC, expression of EGFR and its ligands, transforming growth factor- α or epidermal growth factor, are up-regulated in histologically normal epithelium adjacent to invasive cancer compared with control normal epithelium in individuals without cancer. Dysplastic lesions overexpress EGFR, and invasive cancer displays a more jumped-up pattern of overexpression (19, 20). All this implicates the EGFR signaling pathway in the early stages of head and neck carcinogenesis and progression. EGFR overexpression is also associated with a poorer prognosis in HNC patients (45).

Interaction between EGFR and COX-2 Pathways

Whereas the pathways by which EGFR and COX-2 contribute to carcinogenesis have been separately considered and targeted, increasing evidence indicates a tight connection between these two pathways (Fig. 1).

EGFR and COX-2 signaling pathways form a positive feedback loop. Activation of EGFR has been shown to

induce increased COX-2 expression in various normal and tumor cell lines, including HNSCC cell lines (46–54). Both transforming growth factor- α and epidermal growth factor, ligands of EGFR, were found to induce COX-2 expression (46–50). Expression of COX-2 is regulated at both the transcriptional and posttranscriptional level. The signaling pathway involved in COX-2 induction via EGFR varies depending on the type of cells and inducers, but the ras/raf/mitogen-activated protein kinase signaling pathways mainly contribute to both increased transcriptional and posttranscriptional control. One explanation for the linkage between EGFR/mitogen-activated protein kinase and transcriptional activation of COX-2 may be the activation of transcription factors such as cyclic AMP response element-binding protein/activating transcription factor and activator protein-1 by mitogen-activated protein kinase signaling (25, 46, 51, 54). Binding sites for these transcription activators have been identified in the COX-2 promoter region.

On the other hand, COX-2 induces transactivation or increased expression of EGFR (55, 56). Transactivation of EGFR by PGE₂, a major prostaglandin involved in carcinogenesis, has been well documented but the process seems to be quite complex and cell type dependent. Pai et al. (56) showed that PGE₂ transactivated EGFR and triggered the activation of extracellular signal regulated kinase-2 pathways in normal gastric epithelial cells and colon cancer cells, inducing cell proliferation *in vitro* and *in vivo*. The G-protein coupled receptor, to which the major prostanoids receptors belong, is involved in EGFR transactivation. The mechanism by which G-protein coupled




Figure 1. Interaction between the EGFR and COX-2 pathways. Activation of EGFR induces COX-2 mainly via the mitogen-activated protein kinase pathway. On the other hand, prostaglandins, a product of COX-2, can stimulate EGFR activation. Both EGFR and COX-2 pathways act in common on several aspects of carcinogenesis.

receptor mediates the transactivation of EGFR has not been clearly defined but the release of EGFR ligand by Src-activated transmembrane metalloproteinase has been suggested (31, 57). Consistent with these findings, Pai et al. (56) observed that PGE₂-mediated transactivation of EGFR also involved transforming growth factor- α , likely released by Src-activated metalloproteinase. Complicating these findings, Buchanan et al. (58) reported that the transactivation of EGFR by PGE₂ occurred via an intracellular Src-mediated event but not through the release of an extracellular epidermal growth factor-like ligand in colon cancer cell lines. On the other hand, Shao et al. (34) showed that PGE₂ activated EGFR through the induction of increased amphiregulin expression, one of the EGFR ligands. They showed that PGE₂ activated the cyclic AMP/protein kinase A pathway, which induced expression of amphiregulin in a colon cancer cell line. Induction of EGFR expression by overexpression of COX-1 or COX-2 was reported by Kinoshita et al. (55) in a human colon cancer cell line. There is a recent evidence for EGFR and proliferator-activated receptor transactivation through an Src-dependent pathway (59).

A direct collaborative effect between PGE₂ and EGFR on tumor cell phenotypes, such as invasion and proliferation, is also well documented. Shao et al. showed this collaboration between COX-2/PGE₂ and EGFR pathways (34, 60), observing synergistic induction of amphiregulin by PGE₂ and transforming growth factor- α (34). In a follow-up study, they showed that activation of both PGE₂ and EGFR signaling pathways synergistically promoted the growth and migration of colon cancer cells (60). Pai et al. (61) reported that PGE₂ enhancement of invasiveness in a colon cancer cell line was mediated by transactivation of c-Met-R (hepatocyte growth factor receptor), partly through the transactivation of EGFR. Buchanan et al. (58) made a similar observation, showing that PGE₂-induced cell migration was mediated by the transactivation of EGFR, and associated with intracellular Src activation in colon cancer cells.

It is worth noting that COX inhibitor repressed the EGFR-related pathway and in turn, EGFR inhibitor repressed COX-2 expression, indirectly suggesting their interaction (62, 63). COX inhibitors were reported to block the cell proliferation induced by epidermal growth factor in NIH 3T3 cells, which could be reversed by adding exogenous PGE₂ (62). Gefitinib, an EGFR inhibitor, showed inhibition of COX-2 expression in a HNSCC cell line (63). In addition, celecoxib, a selective COX-2 inhibitor, showed a protective effect against HER-2/neu-induced experimental breast cancer, indirectly suggesting a relationship between the epithelial growth factor receptor family and COX-2 (64).

On top of their known interactions, both the EGFR and COX-2 pathways affect the same aspects of carcinogenesis, such as inhibition of apoptosis and induction of angiogenesis. The conclusion that the EGFR and COX-2 pathways directly or indirectly collaborate in pertinent carcinogenic pathways seems justified.

Therapeutic Implication of Targeting COX-2 and EGFR-Mediated Pathways in Chemoprevention

COX-2 as a Target of Therapeutic and Chemopreventive Agents

Selective COX-2 inhibitors have been developed to avoid interrupting the biosynthesis of prostaglandins by COX-1. Studies in preclinical models have clearly showed that COX-2 inhibitors repressed tumor growth (65). Furthermore, COX inhibitors reduce tumor cell migration and tumor invasiveness as well as inhibit angiogenesis in various cell lines and in a xenograft animal model (66–69). These inhibitory effects have been reported in HNC as well as by *in vivo* and *in vitro* experiments (70–74). In terms of a chemopreventive effect in HNC, Wang et al. (75) reported a significant delay of tumor cell growth and reduced angiogenesis using a COX-2 inhibitor, celecoxib, in a xenograft mouse model. In a head and neck carcinogenesis model, Shiotani et al. (76) pretreated rats with a carcinogen, 4-nitroquinoline-1-oxide, followed by a selective COX-2 inhibitor, nimesulide, at the postinitiation stage. They found that the ingestion of carcinogen induced COX-2 expression in premalignant tongue lesions and squamous cell carcinoma. Subsequent COX-2 inhibitor treatment significantly reduced the development of invasive squamous cell carcinoma. The antitumorigenic properties of these inhibitors are both COX dependent and independent (77, 78).

Based on these promising results, many clinical trials of chemoprevention using various COX-2 inhibitors have been conducted in various organs, although mostly in colon. The first chemopreventive trial using a selective COX-2 inhibitor in humans was conducted on familial adenomatous polyposis patients. Familial adenomatous polyposis patients were treated with celecoxib 400 mg bid for 6 months and were compared with a placebo-treated group for their polyp burden. A significant reduction in polyp burden was observed in the celecoxib-treated group (21). Observing the data from this study, the Food and Drug Administration approved celecoxib as an adjuvant therapy for familial adenomatous polyposis patients. Currently, phase II clinical trials are under way to evaluate COX-2 inhibitors for the prevention of recurrence or development of second primary tumor in early-stage HNC patients and the prevention of cancer in patients with oral leukoplakia or dysplasia, using celecoxib.

However, notable cardiovascular toxicity was reported for specific COX-2 inhibitors recently, which resulted in reevaluation of the clinical use of COX-2 inhibitors (79).³ Studies of COX-2-specific inhibitors, including celecoxib, showed that COX-2 inhibitors increased the thromboembolic cardiovascular risks. A hypothetical explanation is

³ Department of Health and Human Services. NIH halts use of COX-2 inhibitor in large cancer prevention trial. NIH News, December 17, 2004. Available at <http://www.nih.gov/news/pr/dec2004>.

Table 1. Preclinical and clinical studies of combined therapy using COX and EGFR inhibitors

Author (reference)	COX inhibitor	EGFR inhibitor	<i>In vitro/in vivo/clinical</i>
Tortora et al. (94)	SC-236	ZD1839 (plus protein kinase A antisense)	<i>In vitro/vivo</i>
Chen et al. (95)	Celecoxib	ZD1839	<i>In vitro</i>
Zhang et al. (96)	Celecoxib	ZD1839	<i>In vivo</i>
Luca et al. (97)	Refecoxib	ZD1839	<i>In vitro</i>
Torrance et al. (98)	Sulindac	EKB-569	<i>In vivo</i>
Krysan et al. (102)	Celecoxib	OSI-774	<i>In vitro</i>
Reckamp et al. (100)	Celecoxib	ZD1839	Clinical
Writh et al. (101)	Celecoxib	ZD1839	Clinical
Dannenberg et al. (28)	Celecoxib	EKB-569	Clinical*
Choe et al. (this review)	Celecoxib	OSI-774	Clinical*

*HNSCC chemoprevention trials.

that, unlike nonselective nonsteroidal anti-inflammatory drugs, COX-2 inhibitors could not inhibit platelet aggregation (80, 81). Due to concerns about cardiovascular complications associated with long-term use of COX-2 inhibitors, in December 2004, the National Cancer Institute announced the early cessation of a large colorectal cancer prevention clinical trial with celecoxib.³ But in April 2005, the Food and Drug Administration announced restricted use of nonsteroidal anti-inflammatory drugs, including celecoxib, by providing revised labels to include more specific information about the potential cardiovascular and gastrointestinal risks.⁴

EGFR as a Target of Therapeutic or Chemopreventive Agents

A variety of strategies have been developed to block EGFR specifically, including monoclonal antibodies, tyrosine kinase inhibitors, ligand-linked immunotoxins, and antisense approaches (37, 82, 83). Among those strategies, monoclonal antibodies, such as IMC-225 (Cetuximab), and tyrosine kinase inhibitors, such as ZD1839 (Iressa or Gefitinib), and OSI-774 (Tarceva or Erlotinib) have shown promising efficacy and are currently being used in clinical studies singly or in combination with other chemotherapeutic agents or radiotherapy. The antineoplastic effects of EGFR inhibitors include inhibition of cell cycle progression, induction of apoptosis, inhibition of angiogenesis, and decreasing metastasis (40, 84). In HNC, EGFR inhibition also showed growth inhibition and inhibition of metastasis in *in vitro* and *in vivo* experiments (84–89).

Like other anticancer agents, EGFR inhibitor toxicity was also reported. The main toxic effects of EGFR tyrosine kinase inhibitors, such as OSI-774 and ZD1839, include headache, diarrhea, and skin rash (22, 90, 91). Rare association with interstitial lung disease was also observed using ZD1839 (92, 93).

Combined Chemopreventive Therapy Using COX-2 and EGFR Inhibitors

Because the underlying genetic mechanisms of many malignant neoplasms are possibly multipath processes combined with complex cross-talk between pathways, specific blocking of single molecular targets would not outwit the variability and complexity of genetic alterations in cancer. Combined treatments using appropriate multi-agents may be more effective than single agent treatments. In terms of chemoprevention, combining low doses of drugs too toxic for single use may result in negligible toxicity while eluding drug resistance, resulting in an elegant strategy that is both effective and safe.

Considering the tight connection of the COX-2 and EGFR pathways, the combination of their particular inhibitors may block both pathways in a synergistic or additive manner. This idea has been supported by *in vitro* and *in vivo* studies combining COX-2 and EGFR inhibitors (Table 1). Tortora et al. (94) showed a supra-additive inhibitory effect on tumor growth and angiogenesis by combined treatment with SC-236 (COX-2 inhibitor), ZD1839, and protein kinase A antisense in human colon and breast cancer cell lines and a colon cancer xenograft. We also observed a synergistic growth-inhibitory effect by combining celecoxib and ZD1839 in HNSCC *in vitro* (95) and *in vivo* (96). The combination of celecoxib and ZD1839 augmented G₁ cell cycle arrest and further suppressed phosphorylation of EGFR downstream signaling molecules, such as EGFR, extracellular signal regulated kinase, and AKT. This synergistic growth inhibitory effect was also observed in a breast cancer cell line. De Luca et al. (97) found a synergistic growth inhibitory effect on breast cancer cell lines with a combination of rofecoxib and ZD1839. As with our observation, this effect was associated with significant further inactivation of AKT and extracellular signal regulated kinase.

A synergistic growth inhibitory effect also has been achieved in chemoprevention models. Torrance et al. (98) showed that a combination of sulindac, an inhibitor of both COX-1 and COX-2, and EKB-569, an EGFR tyrosine kinase inhibitor, protected APC^{Min/+} mice remarkably

⁴ The Food and Drug Administration announces series of changes to the class of marketed nonsteroidal anti-inflammatory drugs. Food and Drug Administration News, April 7, 2005. Available at <http://www.fda.gov/bbs/topics/news/2005>.

from the development of intestinal neoplasia, compared with the use of single agents alone. Although 100% of the untreated APC^{Min/+} mice developed ~20 polyps, nearly half the mice treated with both agents developed no polyps at all. In HNC, we observed that pretreatment with celecoxib and ZD1839 before tumor injection results in significant tumor growth delay and inhibition compared with control and single-drug-treated groups (96). The same effect could usually be achieved by at least more than double dose of ZD1839 alone (50 mg/kg in the combination versus 150 mg/kg as a single drug; refs. 96, 99). Combining COX-2 and EGFR inhibitors seems to impede or avert several carcinogenic steps with lower toxicity. As observed previously, cellular proliferation is also more efficiently repressed by this combinational strategy. It may inhibit further progression of genetic instability and make abnormal cells more vulnerable to apoptotic cell death. Synergistic inhibition of angiogenesis has also been achieved. It is rational to expect that inhibiting both pathways with a low dose of each inhibitor would reduce toxicity, because the toxicity from COX-2 inhibitor and EGFR inhibitor have unrelated mechanisms.

Based on preclinical studies, several clinical trials using a combination of COX-2 and EGFR inhibitors are either ongoing or pending activation. Two phase I trials using COX-2 and EGFR tyrosine kinase inhibitors were reported in the American Society of Clinical Oncology meetings of last year and this year (100, 101). The first trial enrolled 12 patients with advanced metastatic and recurrent HNSCC (100). Three of nine evaluable patients showed partial response with no dose-limiting toxicities up to the reporting date. The second trial involved 15 patients with advanced NSCLC (101). The combined drugs were well tolerated. Four of 12 evaluable patients showed partial response and three showed stable disease with no unanticipated toxicities. At present, a randomized multicenter trial using celecoxib and EKB-569 (EGFR tyrosine kinase inhibitor) for preventing oral cancer in patients with oral leukoplakia is pending at the M. D. Anderson Cancer Center (28). We are also planning phase I/II chemopreventive clinical trials using a combination of OSI-774 (Tarceva) and celecoxib for former smokers with premalignant lesions and patients with early stage (stage I/II) HNSCC to prevent HNSCC and second primary and recurrent tumors, respectively. Once launched, these clinical trials will provide valuable information on toxicity, response rate, and appropriate biomarkers for using the combination of COX-2 and EGFR inhibitors in cancer clinics.

Future Prospects and Conclusions

In spite of extensive research in EGFR and COX-2 inhibitors, their combined use for chemoprevention is still in its developmental stage. Further investigations are indeed necessary to develop appropriate chemopreventive strategies in HNC. First, the mechanism by which EGFR and COX-2 pathways are deregulated, interact with each other, or contribute to head and neck carcinogenesis must be clarified. Second, more preclinical studies are critical to evaluate the effectiveness of these combinations and to

understand their mechanisms of action as single agents and in combination in HNC specifically. The generation of a good animal model for HNC chemoprevention would be very helpful. Possible drug toxicity from combined use must be evaluated over the long term, although low doses of each drug will be used. Also, substantial biomarkers that can reflect the effect of COX-2 and EGFR combinations need to be identified. Eventually, we may hope to identify the patient population who would derive the most benefit from this combinational chemopreventive approach.

In conclusion, an expanding body of evidence shows tight interaction between the EGFR and COX-2 pathways, which may provide a target for a synergistic inhibitory effect on cancer cell growth using EGFR and COX-2 inhibitors in combination. This combination approach is a promising novel strategy for the chemoprevention of HNC, which may achieve more effective cancer prevention with less drug toxicity or drug resistance than monotherapy with either drug.

References

1. Jemal A, Tiwari RC, Murray T, et al. American Cancer Society. Cancer statistics, 2004. *CA Cancer J Clin* 2004;54:8–29.
2. Vokes EE, Weichselbaum RR, Lippman SM, Hong WK. Head and neck cancer. *N Engl J Med* 1993;328:184–94.
3. Vokes EE, Stenson KM. Therapeutic options for laryngeal cancer. *N Engl J Med* 2003;349:2087–9.
4. Argiris A, Brockstein BE, Haraf DJ, et al. Competing causes of death and second primary tumors in patients with locoregionally advanced head and neck cancer treated with chemoradiotherapy. *Clin Cancer Res* 2004;10:1956–62.
5. Lippman SM, Batsakis JG, Toth BB, et al. Comparison of low-dose isotretinoin with beta carotene to prevent oral carcinogenesis. *N Engl J Med* 1993;328:15–20.
6. Sun SY, Hail N, Lotan R. Apoptosis as a novel target for cancer chemoprevention. *J Natl Cancer Inst* 2004;96:662–72.
7. Gao X, Fisher SG, Mohideen N, Emami B. Second primary cancer patients with laryngeal cancer: a population-based study. *Int J Radiat Oncol Biol Phys* 2003;56:427–35.
8. Slaughter DP, Southwick HW, Smejkal W. Field cancerization in oral stratified squamous epithelium. *Cancer* 1953;6:963–8.
9. Hunter KD, Parkinson EK, Harrison PR. Profiling early head and neck cancer. *Nat Rev Cancer* 2005;5:127–35.
10. O'Shaughnessy JA, Kelloff GJ, Gordon GB, et al. Treatment and prevention of intraepithelial neoplasia: an important target for accelerated new agent development. *Clin Cancer Res* 2002;8:314–46.
11. Papadimitrakopoulou VA, Shin DM, Hong WK. Chemoprevention of head and neck cancer. In: Harrison LB, Sessions RB, Hong WK, editors. *HNC, a multidisciplinary approach*. Philadelphia (PA): Lippincott Williams & Wilkins; 2004. p. 985–1000.
12. Papadimitrakopoulou VA, Clayman GL, Shin DM, et al. Biochemoprevention for dysplastic lesions of the upper aerodigestive tract. *Arch Otolaryngol Head Neck Surg* 1999;125:1083–9.
13. Shin DM, Khuri FR, Murphy B, et al. Combined interferon- α , 13-*cis*-retinoic acid, and α -tocopherol in locally advanced head and neck squamous cell carcinoma: novel bioadjuvant phase II trial. *J Clin Oncol* 2001;19:3010–7.
14. Shin DM, Richards TJ, Seixas-Silva JA, et al. Phase II trial of bioadjuvant therapy with interferon- α 2a (IFN- α 2a), 13-*cis*-retinoic acid (13-*cRA*), and α -tocopherol (α -TF) for locally advanced squamous cell carcinoma of the head and neck (SCCHN): long term follow-up. *Proc Am Soc Clin Oncol* 2003;22:496.
15. Lin DT, Subbaramaiah K, Shah JP, Dannenberg AJ, Boyle JO. Cyclooxygenase-2: a novel molecular target for the prevention and treatment of HNC. *Head Neck* 2002;24:792–9.

16. Ford AC, Grandis JR. Targeting epidermal growth factor receptor in HNC. *Head Neck* 2003;25:67–73.
17. Chan G, Boyle JO, Yang EK, et al. Cyclooxygenase-2 expression is up-regulated in squamous cell carcinoma of the head and neck. *Cancer Res* 1999;9:991–4.
18. Renkonen J, Wolff H, Paavonen T. Expression of cyclooxygenase-2 in human tongue carcinoma and its precursor lesions. *Virchows Arch* 2002;440:594–7.
19. Shin DM, Ro JY, Hong WK, Hittelman WN. Dysregulation of epidermal growth factor receptor expression in premalignant lesions during head and neck tumorigenesis. *Cancer Res* 1994;54:3153–9.
20. Grandis JR, Tweardy DJ, Melhem MF. Asynchronous modulation of transforming growth factor α and epidermal growth factor receptor protein expression in progression of premalignant lesions to head and neck squamous cell carcinoma. *Clin Cancer Res* 1998;4:13–20.
21. Steinbach G, Lynch PM, Phillips RK, et al. The effect of celecoxib, a cyclooxygenase-2 inhibitor, in familial adenomatous polyposis. *N Engl J Med* 2000;342:1946–52.
22. Soulieres D, Senzer NN, Vokes EE, Hidalgo M, Agarwala SS, Siu LL. Multicenter phase II study of erlotinib, an oral epidermal growth factor receptor tyrosine kinase inhibitor, in patients with recurrent or metastatic squamous cell cancer of the head and neck. *J Clin Oncol* 2004;22:77–85.
23. Cohen EE, Rosen F, Stadler WM, et al. Phase II trial of ZD1839 in recurrent or metastatic squamous cell carcinoma of the head and neck. *J Clin Oncol* 2003;21:1980–7.
24. Turini ME, Dubois RN. Cyclooxygenase-2: a therapeutic target. *Annu Rev Med* 2002;53:35–57.
25. Dannenberg AJ, Subbaramaiah K. Targeting cyclooxygenase-2 in human neoplasia: rationale and promise. *Cancer Cell* 2003;4:431–6.
26. Thun MJ, Namboodiri MM, Heath CW, Jr. Aspirin use and reduced risk of fatal colon cancer. *N Engl J Med* 1991;325:1593–6.
27. Oshima M, Dinchuk JE, Kargman SL, et al. Suppression of intestinal polyposis in *Apc*^{Δ716} knockout mice by inhibition of cyclooxygenase 2 (COX-2). *Cell* 1996;87:803–9.
28. Dannenberg AJ, Lippman SM, Mann JR, Subbaramaiah K, DuBois RN. Cyclooxygenase-2 and epidermal growth factor receptor: pharmacologic targets for chemoprevention. *J Clin Oncol* 2005;23:254–66.
29. Choe MS, Rogatko A, Tighiouart M, et al. Role of cyclooxygenase-2 (Cox-2) expression in tumor progression and survival in the squamous cell carcinoma of the head and neck (HNSCC). *Proc Am Soc Clin Oncol* 2005;23:502s.
30. Cao Y, Prescott SM. Many actions of cyclooxygenase-2 in cell dynamics and in cancer. *J Cell Physiol* 2002;190:279–86.
31. Narumiya S, Sugimoto Y, Ushikubi F. Prostanoid receptors: structure, properties, and functions. *Physiol Rev* 1999;79:1193–225.
32. Sheng H, Shao J, Washington MK, Dubois RN. Prostaglandin E₂ increases growth and motility of colorectal carcinoma cells. *J Biol Chem* 2001;276:18075–81.
33. Sonoshita M, Takaku K, Sasaki N, et al. Acceleration of intestinal polyposis through prostaglandin receptor EP2 in *Apc*^{Δ716} knockout mice. *Nat Med* 2001;7:1048–51.
34. Shao J, Lee SB, Guo H, Evers BM, Sheng H. Prostaglandin E₂ stimulates the growth of colon cancer cells via induction of amphiregulin. *Cancer Res* 2003;63:5218–23.
35. Dohadwala M, Batra RK, Luo J, et al. Autocrine/paracrine prostaglandin E₂ production by non-small cell lung cancer cells regulates matrix metalloproteinase-2 and CD44 in cyclooxygenase-2-dependent invasion. *J Biol Chem* 2002;277:50828–33.
36. Eibl G, Reber HA, Hines OJ, Go VLW. COX and PPAR: possible interactions in pancreatic cancer. *Pancreas* 2004;29:247–53.
37. Raymond E, Faivre S, Armand JP. Epidermal growth factor receptor tyrosine kinase as a target for anticancer therapy. *Drugs* 2000;60:15–23.
38. Jorissen RN, Walker F, Puoliot N, Garrett TPJ, Ward CW, Burgess AW. Epidermal growth factor receptor: mechanisms of activation and signaling. *Exp Cell Res* 2003;284:31–53.
39. Holbro T, Civenni G, Hynes NE. The erbB receptors and their role in cancer progression. *Exp Cell Res* 2003;284:99–110.
40. Kari C, Chan TO, Quadros MR, Rodeck U. Targeting the epidermal growth factor receptor in cancer: apoptosis takes center stage. *Cancer Res* 2003;63:1–5.
41. Buettner R, Mora LB, Jove R. Activated STAT signaling in human tumors provides novel molecular targets for therapeutic intervention. *Clin Cancer Res* 2003;8:945–54.
42. Busse D, Doughty RS, Ramsey TT, et al. Reversible G₁ arrest induced by inhibition of the epidermal growth factor receptor tyrosine kinase requires up-regulation of p27^{KIP1} independent of MAPK activity. *J Biol Chem* 2000;275:6987–95.
43. Kong M, Mounier C, Wu J, Posner BI. Epidermal growth factor-induced phosphatidylinositol 3-kinase activation and DNA synthesis: identification of Grb2-associated binder 2 as the major mediator in rat hepatocytes. *J Biol Chem* 2000;275:36035–42.
44. Herbst RS, Shin DM. Monoclonal antibodies to target epidermal growth factor receptor-positive tumors: a new paradigm for cancer therapy. *Cancer* 2002;94:1593–611.
45. Dassonville O, Formento JL, Francoual M, et al. Expression of epidermal growth factor receptor and survival in upper aerodigestive tract cancer. *J Clin Oncol* 1993;11:1873–8.
46. Matsuura H, Sakaue M, Subbaramaiah K, et al. Regulation of cyclooxygenase-2 by interferon γ and transforming growth factor α in normal human epidermal keratinocytes and squamous carcinoma cells: role of mitogen-activated protein kinases. *J Biol Chem* 1999;274:29138–48.
47. Coffey RJ, Hawkey CJ, Damstrup L, et al. Epidermal growth factor receptor activation induces nuclear targeting of cyclooxygenase-2, basolateral release of prostaglandins and mitogenesis in polarizing colon cancer cells. *Proc Natl Acad Sci U S A* 1997;94:657–62.
48. Zakar T, Mijovic JE, Eyster KM, Bhardwaj D, Olson DM. Regulation of prostaglandin H₂ synthase-2 expression in primary human amnion cells by tyrosine kinase dependent mechanisms. *Biochim Biophys Acta* 1998;1391:37–51.
49. Mestre JR, Chan G, Zhang F, et al. Inhibition of cyclooxygenase-2 expression. An approach to preventing HNC. *Ann N Y Acad Sci* 1999;889:62–71.
50. Saha D, Datta PK, Sheng H, et al. Synergistic induction of cyclooxygenase-2 by transforming growth factor- β 1 and epidermal growth factor inhibits apoptosis in epithelial cells. *Neoplasia* 1999;1:508–17.
51. Huh Y, Kim S, Chun J. Differentiation status-dependent regulation of cyclooxygenase-2 expression and prostaglandin E₂ production by epidermal growth factor via mitogen-activated protein kinase in articular chondrocytes. *J Biol Chem* 2003;278:9691–7.
52. Guo Y, Cheng J, Jin G, Gutkind JS, Hellmich MR, Townsend CM, Jr. Gastrin stimulates cyclooxygenase-2 expression in intestinal epithelial cells through multiple signaling pathways: evidence for involvement of ERK5 kinase and transactivation of the epidermal growth factor receptor. *J Biol Chem* 2002;277:48755–63.
53. DuBois RN, Awad J, Morrow J, Roberts LJ, Bishop PR. Regulation of eicosanoid production and mitogenesis in rat intestinal epithelial cells by transforming growth factor- α and phorbol ester. *J Clin Invest* 1994;93:493–8.
54. Chen LC, Chen BK, Chang JM, Chang WC. Essential role of c-Jun induction and coactivator p300 in epidermal growth factor-induced gene expression of cyclooxygenase-2 in human epidermoid carcinoma A431 cells. *Biochim Biophys Acta* 2004;1683:38–48.
55. Kinoshita T, Takahashi Y, Sakashita T, Inoue H, Tanabe T, Yoshimoto T. Growth stimulation and induction of epidermal growth factor receptor by overproduction of cyclooxygenase 1 and 2 in human colon carcinoma cells. *Biochim Biophys Acta* 1999;438:120–30.
56. Pai R, Soreghan B, Szabo IL, Pavelka M, Baatar D, Tarnawski AS. Prostaglandin E₂ transactivates EGF receptor: a novel mechanism for promoting colon cancer growth and gastrointestinal hypertrophy. *Nat Med* 2002;8:289–93.
57. Luttrell LM, Daaka Y, Lefkowitz RJ. Regulation of tyrosine kinase cascades by G-protein coupled receptors. *Curr Opin Cell Biol* 1999;11:177–83.
58. Buchanan FG, Wang D, Bargiacchi F, DuBois RN. Prostaglandin E₂ regulates cell migration via the intracellular activation of the epidermal growth factor receptor. *J Biol Chem* 2003;278:35451–7.
59. Slomiany BL, Slomiany A. Role of epidermal growth factor transactivation in PPAR[γ]-dependent suppression of helicobacter pylori interference with gastric mucin synthesis. *Inflammopharmacology* 2004;12:177–88.
60. Shao J, Evers BM, Sheng H. Prostaglandin E₂ synergistically enhances receptor tyrosine kinase-dependent signaling system in colon cancer cells. *J Biol Chem* 2004;279:14287–93.

61. Pai R, Nakamura T, Moon WS, Tarnawski AS. Prostaglandins promote colon cancer cell invasion; signaling by cross-talk between two distinct growth factor receptors. *FASEB J* 2003;17:1640–7.
62. Handler JA, Danilowicz RM, Eling TE. Mitogenic signaling by epidermal growth factor (EGF), but not platelet-derived growth factor, requires arachnoid acid metabolism in BALB/c 3T3 cells: modulation of EGF-dependent c-myc expression by prostaglandins. *J Biol Chem* 1990; 265:3669–73.
63. De Lorenzo MS, Chang M, Du B, et al. Gefitinib ('Iressa', ZD1839) inhibits the expression of Cox-2 and the growth of experimental head and neck squamous-cell carcinoma. *Proc Am Assoc Cancer Res* 2004;45:1072.
64. Howe LR, Subbaramaiah K, Patel J, et al. Celecoxib, a selective cyclooxygenase 2 inhibitor, protects against human epidermal growth factor receptor 2 (HER-2)/neu-induced breast cancer. *Cancer Res* 2002; 62:5405–7.
65. Subbaramaiah K, Dannenberg AJ. Cyclooxygenase 2: a molecular target for cancer prevention and treatment. *Trends Pharmacol Sci* 2003;24:96–102.
66. Attiga FA, Fernandez PM, Weeraratna AT, Manyak MJ, Patierno SR. Inhibitors of prostaglandin synthesis inhibit human prostate tumor cell invasiveness and reduce the release of matrix metalloproteinases. *Cancer Res* 2000;60:4629–37.
67. Chen W, Wei S, Liu JM, Hsiao M, Kou-Lin J, Yang WK. Tumor invasiveness and liver metastasis of colon cancer cells correlates with cyclooxygenase-2 (COX-2) expression and inhibited by a COX-2-selective inhibitor, etodolac. *Int J Cancer* 2001;91:894–9.
68. Rozic JG, Chakraborty C, Lala PK. Cyclooxygenase inhibitors retard murine mammary tumor progression by reducing tumor cell migration, invasiveness and angiogenesis. *Int J Cancer* 2001;93:497–506.
69. Masferrer JL, Leahy KM, Koki AT, et al. Antiangiogenic and antitumor activities of cyclooxygenase-2 inhibitors. *Cancer Res* 2000;60:1306–11.
70. Peng JP, Liu LT, Chang HC, Hung WC. Enhancement of chemotherapeutic drug-induced apoptosis by a cyclooxygenase-2 inhibitor in hypopharyngeal carcinoma cells. *Cancer Lett* 2003;201:157–63.
71. Hashitani S, Urade M, Nishimura N, et al. Apoptosis induction and enhancement of cytotoxicity of anticancer drugs by celecoxib, a selective cyclooxygenase-2 inhibitor, in human head and neck carcinoma cell lines. *Int J Oncol* 2003;23:665–72.
72. Minte HA, Eveson JW, Huntley S, Elder DJ, Hague A. The cyclooxygenase 2-selective inhibitor NS398 inhibits proliferation of oral carcinoma cell lines by mechanisms dependent and independent of reduced prostaglandin E₂ synthesis. *Clin Cancer Res* 2003;9: 1885–97.
73. Zweifel BS, Davis TW, Ornberg RL, Masferrer JL. Direct evidence for a role of cyclooxygenase 2-derived prostaglandin E₂ in human head and neck xenograft tumors. *Cancer Res* 2002;62:6706–11.
74. Sumitani K, Kamijo R, Toyoshima T, et al. Specific inhibition of cyclooxygenase-2 results in inhibition of proliferation of oral cancer cell lines via suppression of prostaglandin E₂ production. *J Oral Pathol Med* 2001;30:41–7.
75. Wang Z, Fuentes CF, Shapshay SM. Antiangiogenic and chemopreventive activities of celecoxib in oral carcinoma cell. *Laryngoscope* 2002;112:839–43.
76. Shiotani H, Denda A, Yamamoto K, et al. Increased expression of cyclooxygenase-2 protein in 4-nitroquinoline-1-oxide-induced rat tongue carcinomas and chemopreventive efficacy of a specific inhibitor, nimesulide. *Cancer Res* 2001;61:1451–6.
77. Hwang DH, Fung V, Dannenberg AJ. National cancer institute workshop on chemopreventive properties of nonsteroidal anti-inflammatory drugs: role of cox-dependent and -independent mechanisms. *Neoplasia* 2002;4:91–7.
78. Song X, Lin H, Johnson AJ, et al. Cyclooxygenase-2, player or spectator in cyclooxygenase-2 inhibitor-induced apoptosis in prostatic cancer cells. *J Natl Cancer Inst* 2002;94:585–91.
79. Bannwarth B. Do selective cyclo-oxygenase-2 inhibitors have a future? *Drug Safety* 2005;28:183–9.
80. FitzGerald GA. Coxibs and cardiovascular disease. *N Engl J Med* 2004;351:1709–11.
81. White WB, Faich G, Borer JS, Makuch RW. Cardiovascular thrombotic events in arthritis trials of the cyclooxygenase-2 inhibitors celecoxib. *Am J Cardiol* 2003;92:411–8.
82. Khalil MY, Grandis JR, Shin DM. Targeting epidermal growth factor receptor: novel therapeutics in the management of cancer. *Expert Rev Anticancer Ther* 2003;3:367–80.
83. Grunwald V, Hidalgo M. Developing inhibitors of the epidermal growth factor receptor for cancer treatment. *J Natl Cancer Inst* 2003;95:851–67.
84. Matheny KE, Barbieri CE, Sniezek JC, Arteaga CL, Pietenpol JA. Inhibition of epidermal growth factor receptor signaling decreases p63 expression in head and neck squamous carcinoma cells. *Laryngoscope* 2003;113:936–9.
85. Di Gennaro E, Barbarino M, Bruzzese F, et al. Critical role of both p27KIP1 and p21CIP1/WAF1 in the antiproliferative effect of ZD1839 ('Iressa'), an epidermal growth factor receptor tyrosine kinase inhibitor, in head and neck squamous carcinoma cells. *J Cell Physiol* 2003;195: 139–50.
86. Shintani S, Li C, Mihara M, et al. Gefitinib ('Iressa', ZD1839), an epidermal growth factor receptor tyrosine kinase inhibitor, up-regulates p27KIP1 and induces G₁ arrest in oral squamous cell carcinoma cell lines. *Oral Oncol* 2004;40:43–51.
87. Shintani S, Li C, Mihara M, Nakashiro K, Hamakawa H. Gefitinib ('Iressa'), an epidermal growth factor receptor tyrosine kinase inhibitor, mediates the inhibition of lymph node metastasis in oral cancer cells. *Cancer Lett* 2003;201:149–55.
88. Faust RA, Tawfic S, Davis AT, Ahmed K. Apoptosis and growth inhibition of head and neck tumor cell line induced by epidermal growth factor receptor tyrosine kinase inhibitor. *Oral Oncol* 1999;35:290–5.
89. Modjtahedi H, Affleck K, Stubberfield C, Dean C. EGFR blockade by tyrosine kinase inhibitor or monoclonal antibody inhibits growth, directs terminal differentiation and induces apoptosis in the human squamous cell carcinoma HN5. *Int J Oncol* 1998;13:335–42.
90. Barlesi FA, Tchouhadjian CA, Daddoli C, et al. Gefitinib [(ZD1839), Iressa (R)] in non-small-cell lung cancer: a review of clinical trials from daily practice perspective. *Fund Clin Pharm* 2005;19:385–93.
91. Hidalgo M, Siu LL, Nemunaitis J, et al. Phase I and pharmacologic study of OSI-774, an epidermal growth factor receptor tyrosine kinase inhibitor, in patients with advanced solid malignancies. *J Clin Oncol* 2001; 19:3267–79.
92. Inoue A, Saijo Y, Maemondo M, et al. Severe acute interstitial pneumonia and gefitinib. *Lancet* 2003;361:137–9.
93. Forsythe B, Faulkner K. Overview of the tolerability of gefitinib (IRESSA[®]) monotherapy. *Drug Safety* 2004;27:1081–92.
94. Tortora G, Caputo R, Damiano V, et al. Combination of a selective cyclooxygenase-2 inhibitor with epidermal growth factor receptor tyrosine kinase inhibitor ZD1839 and protein kinase A antisense causes cooperative antitumor and antiangiogenic effect. *Clin Cancer Res* 2003;9:1566–72.
95. Chen Z, Zhang X, Li M, et al. Simultaneously targeting epidermal growth factor receptor tyrosine kinase and cyclooxygenase-2, an efficient approach to inhibition of squamous cell carcinoma of the head and neck. *Clin Cancer Res* 2004;10:5930–9.
96. Zhang X, Chen Z, Choe MS, et al. Tumor growth inhibition by simultaneously blocking EGFR a cyclooxygenase-2 in a xenograft model. *Clin Cancer Res*. In press. 2005.
97. De Luca A, Maiello MR, Perrone F, et al. Cooperative inhibitory effect of gefitinib ('Iressa', ZD1839) in combination with rofecoxib on human breast cancer cell. *Proc Am Assoc Cancer Res* 2004;45:894.
98. Torrance CJ, Jackson PE, Montgomery E, et al. Combinatorial chemoprevention of intestinal neoplasia. *Nat Med* 2000;6:1024–8.
99. Sirotnak FM, Zakowski MF, Miller VA, Scher HI, Kris MG. Efficacy of cytotoxic agents against human tumor xenografts is markedly enhanced by coadministration of ZD1839 (Iressa), an inhibitor of EGFR tyrosine kinase. *Clin Cancer Res* 2000;6:4885–92.
100. Writh LJ, Haddad RI, Wiczorek TJ, et al. Phase I study of gefitinib plus celecoxib in patients with metastatic and/or locally recurrent squamous cell carcinoma of the head and neck (SCCHN). *Proc Am Soc Clin Oncol* 2004;40:496.
101. Reckamp KL, Dubinett SM, Krysan K, Figlin RA. A phase I trial of targeted COX-2 and EGFR TK inhibition in advanced NSCLC. *Proc Am Soc Clin Oncol* 2005;41:648s.
102. Krysan K, Reckamp KL, Escudero B, Bando JM, Dubinett SM. PGE₂-stimulated proliferation of NSCLC cells is epidermal growth factor receptor-independent and resistant to erlotinib. *Proc Am Assoc Cancer Res* 2005;46:36.

Molecular Cancer Therapeutics

Interaction between epidermal growth factor receptor– and cyclooxygenase 2–mediated pathways and its implications for the chemoprevention of head and neck cancer

Mi Sun Choe, Xin Zhang, Hyung Ju C. Shin, et al.

Mol Cancer Ther 2005;4:1448-1455.

Updated version Access the most recent version of this article at:
<http://mct.aacrjournals.org/content/4/9/1448>

Cited articles This article cites 98 articles, 37 of which you can access for free at:
<http://mct.aacrjournals.org/content/4/9/1448.full.html#ref-list-1>

Citing articles This article has been cited by 17 HighWire-hosted articles. Access the articles at:
<http://mct.aacrjournals.org/content/4/9/1448.full.html#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.