

**Article category**

Original article

**Article title**

Clinical features and prognostic factors of early-onset sepsis : a 7.5-year experience in one neonatal intensive care unit

**Running title**

Clinical features and prognostic factors of early-onset sepsis

**Author and Institution:**

Se Jin Kim, Ga Eun Kim, Jae Hyun Park, Sang Lak Lee, and Chun Soo Kim

Department of Pediatrics, Keimyung University School of Medicine, Daegu, Korea

**Information of Corresponding author:**

Chun Soo Kim, MD, PhD

Department of Pediatrics, Keimyung University School of Medicine, 56 Dalseong-ro, Jung-gu, Daegu 41931, Korea

Tel: +82-53-250-7526

Fax: +82-53- 250-7783

E-mail: cskim@dsmc.or.kr

## Abstract

**Purpose:** This study was undertaken to investigate the clinical features and prognostic factors of early-onset sepsis (EOS) in neonatal intensive care unit (NICU) patients.

**Methods:** A retrospective analysis of the medical records was conducted in a NICU of a university hospital over a 7.5-year period (Jan 2010-Jun 2017).

**Results:** During the study period, there were 45 (1.2%) episodes of EOS in 3,862 infants. The most common pathogen responsible for EOS was *Streptococcus* group B (GBS) in 10 cases (22.2%), followed by *E. coli* in 9 cases (20%). The frequency of Gram-positive sepsis was higher in term than in preterm infants, while the rate of Gram-negative infection was higher in preterm than in term infants ( $P<0.05$ ). The overall mortality was 37.8% (17 of 45), and 47% of deaths occurred within the first 3 days of infection. There was a significant difference in terms of gestational age (26.8 weeks vs. 35.1 weeks) and birth weight (957 g vs. 2,520 g) between the death and survival groups. In a comparison after adjusting for the difference in gestational age and birth weight between the two groups, Gram-negative pathogens (OR: 42, 95% CI: 1.4~1281.8) and some clinical findings, such as neutropenia (OR: 46, 95% CI: 1.3~1628.7) and decreased activity (OR: 34, 95% CI: 1.8~633.4), were associated with fatality.

**Conclusion:** The common pathogens responsible for EOS in NICU patients are GBS and *E. coli*. Infection caused by Gram-negative bacteria, decreased activity in the early phase of infection, and neutropenia were associated with poor outcomes.

**Key words:** Early-onset sepsis; Clinical features, Prognosis, Neonatal intensive care unit

## Introduction

Although the survival rate of newborn infants has been improved by advance in neonatal intensive therapy, the incidence of infections in high-risk infants is not decreasing. The overall incidence of early-onset sepsis (EOS), defined as sepsis occurring within 72 hours of birth, is 1-4.6 per 1,000 live births worldwide,<sup>1,2)</sup> but increases to 11-36 per 1,000 live births in very low birth weight infants (VLBWIs).<sup>1,3,4)</sup> VLBWIs with EOS have a higher rate of mortality and complications, including bronchopulmonary dysplasia, severe ventricular hemorrhage, sever retinopathy of prematurity and neurodevelopmental defects, than those without infection.<sup>3-5)</sup>

EOS is usually caused by microorganisms, including bacteria, that are present in the birth canal of the mother, and the infection occurs before or during labor. Various obstetric factors, including chorioamnionitis, premature rupture of membranes (PROM), maternal fever, and preterm labor, are associated with EOS.<sup>6-8)</sup> The incidence of EOS and the distribution of causative organisms vary by geographic region and race,<sup>9,10)</sup> and also depends on the medical environment at individual hospitals, such as the application of maternal chemoprophylaxis and the proportion of VLBWI among neonatal intensive care unit (NICU) patients.<sup>9,11)</sup> Intrapartum antibiotic prophylaxis (IAP) using ampicillin has been administered to pregnant women with group B *Streptococcus* (GBS) rectovaginal colonization and other obstetric risk factors for EOS in the United States (US) since 1996, and the incidence of EOS due to GBS has decreased.<sup>12)</sup> However, infection by *E. coli* has not been reduced,<sup>13)</sup> and infection by resistant strains has been increasing.<sup>14)</sup> Therefore, serial investigations of the causative organisms of EOS in individual NICUs are important for infection control.

Infants suffering from EOS show a high mortality rate of approximately 11-37%,<sup>1,3,13)</sup> despite advance in intensive care, including early antibiotic therapy. The time of death due to

sepsis is usually within 2 weeks after the onset of the infection.<sup>1,5)</sup> The incidence of fatality is affected by the gestational age and birth weight of babies with infections.<sup>1)</sup> Additionally, the outcomes of sepsis may differ depending on the causative pathogens and clinical findings for the infected patients.<sup>13,15,16)</sup> However, clinical studies on the prognostic factors associated with the mortality of EOS are still lacking. This study was undertaken to investigate the clinical features and prognostic factors related to fatality of EOS in NICU patients.

## **Materials and methods**

This study was conducted on patients with EOS at the NICU of a university hospital in Korea over a 7.5-year period (Jan 2010-Jun 2017). The medical records and radiographic findings of the subjects were retrospectively reviewed. This study was approved by the institutional review board of Keimyung University Dongsan Hospital (IRB No. 2018-06-027). Informed consent was waived by the board.

### **1. Study design**

The subjects were those who had clinical signs of infection and/or elevated C-reactive protein (CRP) among the patients who were identified with causative organisms in blood cultures within 72 hours of life.<sup>1,3)</sup> In patients with skin flora, including coagulase-negative *staphylococci* (CONS), the diagnosis of sepsis was based on both the presence of clinical signs of infection and elevated CRP, or the same bacteria were continuously detected in the blood culture performed within 48 hours. Among patients who were positive for microorganisms in their blood specimens within 3 days of life, those with contaminants were

excluded from the study. The following were assessed as contamination: if contaminants (*Corynebacterium*, *Diphtheroids*, *Micrococci*, and *Propionibacterium*) or more than one organism were detected in the blood culture and antibiotics were discontinued within 5 days in the surviving patient.<sup>1,5)</sup> Using the medical data of patients with EOS, the incidence of sepsis and clinical findings of infections, including causative organisms, symptoms and signs, laboratory findings, mortality, and risk factors associated with fatality, were investigated.

## 2. Definitions of variables

Clinical symptoms associated with sepsis were limited to the following: decreased activity, fever, tachycardia, bradycardia, apnea, abdominal distension, and hypotension. Clinical symptoms and laboratory tests in the early phase of infection were limited to findings within the first 48 hours from the time of blood culture. Chorioamnionitis included both a clinical and histological diagnosis.<sup>8)</sup> Maternal hypertension included all four types of hypertension in pregnancy classified as gestational hypertension, preeclampsia, eclampsia, and chronic hypertension of any etiology.<sup>17)</sup> The use of antenatal steroids was limited to 2 or more doses of dexamethasone. The use of antenatal antibiotics was limited to the administration of drugs for more than 3 days. Apnea was defined as temporary cessation of breathing for a duration of >20 seconds or when accompanied by bradycardia and desaturation, even though the duration was <20 seconds.<sup>18)</sup> Hypotension was defined as a mean blood pressure of <30 mmHg.<sup>19)</sup> Neutropenia was defined as a neutrophil count of <1,000/ $\mu$ L.<sup>20)</sup> Respiratory distress syndrome (RDS) was limited to the cases undergoing surfactant replacement treatments. Pneumothorax was limited to cases with chest tube insertion. Pulmonary hemorrhage was limited to cases that required interventional therapies, including tracheal

instillation of epinephrine. Patent ductus arteriosus (PDA) was limited to cases with hemodynamic significance confirmed by echocardiography and associated symptoms. Acute kidney injury was defined as both oliguria (<1 mL/kg/h) and elevated serum creatinine (>1.5 mg/dL).<sup>21)</sup> Empirical antimicrobial therapy for EOS was based on the use of ampicillin and gentamicin. Also, the results of culture test for maternal specimens were considered. The empirical antimicrobial regimen was considered appropriate if at least one antibiotic is susceptible to the causative organism; otherwise, it was assessed to be inappropriate. Fulminant infection was defined as deaths within 48 hours after the onset of sepsis.<sup>22)</sup> Fatality in patients with sepsis was limited to death within 14 days after infection.

### **3. Statistical analysis**

Data were analyzed using SPSS Statistics version 20.0 (Apache Software Foundation, Chicago, IL, USA). The chi-square test and Fisher's exact test were used for categorical variables, and the Man-Whitney U test and t test were used for continuous variables. Multiple logistic regression analysis was performed to detect the risk factors associated with the prognosis of EOS. P-values less than 0.05 were considered statistically significant.

## **Results**

### **1. Study population**

Of the total 3,862 patients, microorganisms were identified in the blood samples of 167 babies (4.3%). Among the infants with identified microbes in their blood specimens, 45 were

diagnosed as infected, and the remaining 122 were classified as cases with contaminants. Among the babies with contaminants, mixed bacteria were detected in 2 cases, and 2 different contaminants were detected in blood culture tests at intervals of 48 hours in another. The EOS group was classified into 17 deaths and 28 survivors (Fig. 1).

## **2. Distribution of microorganisms**

In a total of 45 subjects, the most common pathogen responsible for EOS was GBS in 10 cases (22.2%), followed by *E. coli* in 9 cases (20%). The rate of Gram-positive sepsis was higher in term than in preterm infants (85.7% vs. 51.6%) ( $P<0.05$ ). In contrast, the rate of Gram-negative infection was higher in preterm than in term infants (45.2% vs. 14.3%) ( $P<0.05$ ) (Table 1).

## **3. Mortality and time of death**

The overall mortality within 14 days after the infection was 37.8% (17 of 45). The number of deaths was the highest at 8 (47.1%) during the first 3 days after infection, 4 (23.5%) after 4-7 days, and 5 (29.4%) after 8-14 days (Table 2). Fulminant infections occurred in 7 cases (15.6%), and all were preterm infants. In detail, 3 cases occurred in patients with *P. aeruginosa* and *E. coli* sepsis, respectively; the remaining 1 developed in infants with *S. aureus* sepsis.

## **4. Incidence of fatal infections by causative organisms**

The rate of fatality in patients with *P. aeruginosa*, *E. coli* and GBS sepsis was 100%, 77.8%, and 10%, respectively. The overall rate of fatal infections in infants with Gram-negative sepsis was 68.8%, which was higher than the 21.4% for patients with Gram-positive infections ( $P<0.005$ ) (Table 3).

## 5. Perinatal and clinical findings

There was a significant difference in terms of gestational age ( $26.8\pm 3.3$  weeks vs.  $35.1\pm 4.3$  weeks) and birth weight ( $957.1\pm 542.4$  g vs.  $2,520.4\pm 924.4$  g) between the death and survival groups ( $P<0.001$ ). Furthermore, the proportion of VLBWI was higher in the death group than in the survival group (82.4% vs. 14.3%) ( $P<0.001$ ). The 1-min and 5-min Apgar scores were lower in the death group ( $P<0.001$ ). The frequency of chorioamnionitis, antenatal steroid and antibiotic therapy was also higher in the death group ( $P<0.05$ ). In a comparison of clinical symptoms during the early phase of infection, the frequencies of several variables, including decreased activity, bradycardia, hypotension, and fever, were different between the death and survival groups ( $P<0.05$ ). Among the laboratory findings, elevated CRP and leukopenia were the most common findings (64.4% and 33.3%, respectively); however, there were no differences between the two groups. The rate of neutropenia was significantly higher in the death group than in the survival group ( $P<0.005$ ). Moreover, in a comparison of comorbid diseases in the early phase of infection, several factors, including RDS, PDA, and pulmonary hemorrhage, were more common in the death group than in the survival group ( $P<0.05$ ). Meningitis was complicated in 2 term infants with GBS sepsis, both of whom survived. The appropriateness of empiric antimicrobial therapy was not different between the two groups (Table 4).



## 6. Prognostic factors of fatal outcome

Logistic regression analysis of individual variables was performed to adjust for differences in gestational age and birth weight between the death and survival groups. Gram-negative pathogens (OR: 42, 95% CI: 1.4~1281.8) and some clinical findings, such as neutropenia (OR: 46, 95% CI: 1.3~1628.7) and decreased activity (OR: 34, 95% CI: 1.8~633.4), were associated with fatality of sepsis after adjusting for these differences (Table 5).

## Discussion

According to recent data from the Korean Neonatal Network (KNN),<sup>4)</sup> the incidence of EOS, defined as sepsis occurring within 3 days of life, was 3.6%. In this study, according to data from a single hospital over a 7.5-year period, EOS developed in 1.2% of all NICU patients. The KNN investigated VLBWI among NICU patients. The difference between the incidence reported by the KNN and this study can be explained by the differences in gestational age and birth weight of the infants in the 2 study populations. The incidence of EOS increases with lower gestational age and body weight. In a national study performed in the US,<sup>1)</sup> the overall rate of EOS was 0.98 cases per 1,000 live births. Specifically, rates of infection were 10.96, 1.38, and 0.57 per 1,000 live births in infants with a birth weight of 401-1,500 g, 1,501-2,500 g, and >2,500 g, respectively.

Generally, pathogens causing EOS are more common in Gram-positive bacteria than Gram-negative bacteria,<sup>1,9,13)</sup> and the most frequent pathogen is GBS.<sup>1,9)</sup> However, according to data limited to VLBWI, Gram-negative pathogens are more common than Gram-positive

organisms,<sup>3)</sup> and *E. coli* is the most common pathogen.<sup>1,3)</sup> In developing countries where the level of health care is low and the delivery environment is poorly sanitized, other Gram-negative bacteria, including *K. pneumonia* and *E. cloacae*, may be the most common pathogens.<sup>10)</sup> The incidence of EOS by GBS infection has been decreasing since the introduction of IAP in the US, with 1.7 cases per 1,000 live births in the early 1990s and 0.34-0.37 cases per 1,000 live births in 2003-2008 period.<sup>12)</sup> Nonetheless, GBS is still the most common organism in neonatal EOS, while *E. coli* is an increasing pathogen, especially in preterm infants.<sup>13)</sup> In this study, similar to previous reports, the common pathogens responsible for EOS were GBS and *E. coli*. In detail, the number of cases with GBS sepsis was the same in both term and preterm infants (5 vs. 5); however, the number of cases with *E. coli* infection was higher in preterm than in full-term infants (8 vs. 1).

The mortality of neonatal EOS is inversely related to gestational age and birth weight.<sup>1)</sup> In the current study of total NICU patients regardless of gestational age, the mortality rate for patients with EOS was 37.8%, which can be considered as high. However, the fact that the mean gestational age was 27.8 weeks and the mean birth weight was 957.1 g in the death group should be considered, and direct comparison is difficult because the demographic characteristics of the subjects were different in each study.<sup>1,4)</sup> Moreover, the mortality rate of infants with sepsis varies according to causative pathogens,<sup>9,15)</sup> and Gram-negative infection is associated with a higher mortality than Gram-positive infection.<sup>13,15)</sup> In this study, the mortality of patients with Gram-negative sepsis was 68.8%, which was higher than the 21.4% of patients with Gram-positive sepsis. Specifically, the mortality of infants with *E. coli* sepsis was 77.8%, while the mortality of infants with GBS sepsis was 10%. Fulminant infections that result in fatal outcomes in the early phase of infections are often associated with Gram-negative sepsis, including *P. aeruginosa* and *E. coli*, especially in extremely preterm infants

with septic shock.<sup>16,22)</sup> In this study, 6 of the 7 fulminant infections were caused by Gram-negative sepsis, and 5 of the affected patients were VLBWI. The shorter the time interval between the onset of infection and the time of death in patients with sepsis, the more likely it is that the infection was directly responsible for the death.<sup>23)</sup> In this study, the proportions of deaths during the first 3 days and first 7 days after the onset of infection were 47.1% and 70.6%, respectively, similar to previous reports.<sup>1,5)</sup>

The three major risk factors for EOS in newborn infants are chorioamnionitis, PROM  $\geq 18$  hours, and inadequate IAP in the at-risk group.<sup>24)</sup> In infants with EOS, the rate of chorioamnionitis is 33-77%,<sup>1,4,5,8)</sup> and this varies depending on the diagnostic method or the compliance with histologic examination. In this study, both clinical and histologic findings were included in the diagnosis, and the rate of chorioamnionitis was 51.1%, however, histologic examinations were not performed in some patients. Although 33-44% of patients with EOS are reported to have prolonged PROM,<sup>1,4,5)</sup> only 13.3% of the subjects in this study were affected. The application of IAP in pregnant women with risk factors, including rectovaginal colonization of GBS, PROM, fever, or preterm labor, is effective in preventing GBS sepsis;<sup>12)</sup> however, the guideline are yet to be established in Korea. IAP has recently been applied to some patients during this study period. Therefore, it is necessary to evaluate the effectiveness of chemoprophylaxis in the future. The rate of concomitant meningitis in infants with EOS is reported to be 4-9%,<sup>1,13,25,26)</sup> and it developed in 4.4% of the subjects in this study. Concomitant meningitis can be a poor prognostic factor in infants with EOS; specifically, meningitis caused by Gram-negative bacteria in VLBWI is associated with fatal outcomes.<sup>25)</sup> In this study, meningitis occurred in 2 term infants with GBS sepsis but was not associated with death.

In this study, logistic regression analysis was performed to adjust for the significant

differences in gestational age (26.8 weeks vs. 35.1 weeks) and birth weight (957 g vs. 2,520 g) between the death and survival groups. Gram-negative pathogens (OR: 42, 95% CI: 1.4~1281.8) and some clinical findings, such as neutropenia (OR: 46, 95% CI: 1.3~1628.7) and decreased activity (OR: 34, 95% CI: 1.8~633.4), were associated with fatal outcome of sepsis after adjusting for these differences. The appearance of neutropenia in the peripheral blood of newborn infants with sepsis suggests that the neutrophil storage pool is depleted.<sup>27)</sup> Sepsis in neutropenic patients often presents as severe infections, and it can rapidly progress to septic shock and multiple organ dysfunction syndromes and is associated with high mortality.<sup>28,29)</sup> In addition, decreased activity in babies with sepsis can be considered as a clinical symptom related to acute changes in mental status, suggesting severe infection.<sup>30)</sup> Therefore, decreased activity seen in the early phase of infection in infants with sepsis, may be associated with a poor prognosis.<sup>31)</sup> This study had some limitations. First, it was retrospective and investigated limited clinical findings that were comparable between the death and survival groups in infants with EOS. Furthermore, it was carried out at a single institution, and the number of subjects was relatively small. However, the results of this study are helpful to understand the clinical features and prognostic factors of EOS. In conclusion, the common pathogens involved in EOS in NICU patients were GBS and *E. coli*. Moreover, infection caused by Gram-negative bacteria, decreased activity in the early phase of infection, and neutropenia are associated with poor outcomes.

### **Conflict of Interest**

The authors have no conflicts of interest to disclose.

## References

1. Stoll BJ, Hansen NI, Sánchez PJ, Faix RG, Poindexter BB, Van Meurs KP, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Early onset neonatal sepsis: the burden of group B Streptococcal and *E. coli* disease continues. *Pediatrics* 2011;127:817-26.
2. Edwards RK, Jamie WE, Sterner D, Gentry S, Counts K, Duff P. Intrapartum antibiotic prophylaxis and early-onset neonatal sepsis patterns. *Infect Dis Obstet Gynecol* 2003;11:221-6.
3. Stoll BJ, Hansen N, Fanaroff AA, Wright LL, Carlo WA, Ehrenkranz RA, et al. Changes in pathogens causing early-onset sepsis in very-low-birth-weight infants. *N Engl J Med* 2002;347:240-7.
4. Lee SM, Chang M, Kim KS. Blood culture proven early onset sepsis and late onset sepsis in very-low-birth-weight infants in Korea. *J Korean Med Sci* 2015;Suppl 1:S67-74.
5. Klinger G, Levy I, Sirota L, Boyko V, Lerner-Geva L, Reichman B; Israel Neonatal Network. Outcome of early-onset sepsis in a national cohort of very low birth weight infants. *Pediatrics* 2010;125:e736-40.
6. Schuchat A, Zywicki SS, Dinsmoor MJ, Mercer B, Romaguera J, O'Sullivan MJ, et al. Risk factors and opportunities for prevention of early-onset neonatal sepsis: a multicenter case-control study. *Pediatrics* 2000;105:21-6.
7. Kim DH, Lee HJ, Kim HS, Yoo BH. Influence of histologic chorioamnionitis and funisitis on the level of peripheral blood C-reactive protein at birth in preterm infants. *Korean J Pediatr* 2010;53:33-40.
8. Lee SY, Park KH, Jeong EH, Oh KJ, Ryu A, Park KU. Relationship between maternal

- serum C-reactive protein, funisitis and early-onset neonatal sepsis. *J Korean Med Sci* 2012;27:674-80.
9. Simonsen KA, Anderson-Berry AL, Delair SF, Davies HD. Early-onset neonatal sepsis. *Clin Microbiol Rev* 2014;27:21-47.
  10. Fahmey SS. Early-onset sepsis in a neonatal intensive care unit in Beni Suef, Egypt: bacterial isolates and antibiotic resistance pattern. *Korean J Pediatr* 2013;56:332-7.
  11. Shane AL, Stoll BJ. Neonatal sepsis: progress towards improved outcomes. *J Infect* 2014;68 Suppl 1:S24-32.
  12. Verani JR, McGee L, Schrag SJ. Prevention of perinatal group B streptococcal disease--revised guidelines from CDC, 2010. *MMWR Recomm Rep* 2010;59:1-36.
  13. Schrag SJ, Farley MM, Petit S, Reingold A, Weston EJ, Pondo T, et al. Epidemiology of invasive early-onset neonatal sepsis, 2005 to 2014. *Pediatrics* 2016;138:e20162013.
  14. Wright AJ, Unger S, Coleman BL, Lam PP, McGeer AJ. Maternal antibiotic exposure and risk of antibiotic resistance in neonatal early-onset sepsis: a case-cohort study. *Pediatr Infect Dis J* 2012;31:1206-8.
  15. Piening BC, Geffers C, Gastmeier P, Schwab F. Pathogen-specific mortality in very low birth weight infants with primary bloodstream infection. *PLoS One* 2017;12:e0180134.
  16. Kermorvant-Duchemin E, Laborie S, Rabilloud M, Lapillonne A, Claris O. Outcome and prognostic factors in neonates with septic shock. *Pediatr Crit Care Med* 2008;9:186-91.
  17. Koo YJ, Lee DH. Hypertension in pregnancy. *J Korean Med Assoc* 2016;59:24-30.
  18. Sale SM. Neonatal apnoea. *Best Pract Res Clin Anaesthesiol* 2010;24:323-36.
  19. Park HK. Optimal Blood Pressure in Preterm Infants. *Neonatal Med* 2014;21:99-105.
  20. Nittala S, Subbarao GC, Maheshwari A. Evaluation of neutropenia and neutrophilia in preterm infants. *J Matern Fetal Neonatal Med* 2012;25(Suppl 5):100-3. doi:

10.3109/14767058.2012.715468.

21. Subramanian S, Agarwal R, Deorari AK, Paul VK, Bagga A. Acute renal failure in neonates. *Indian J Pediatr* 2008;75:385-91.
22. Karlowicz MG, Buescher ES, Surka AE. Fulminant late-onset sepsis in a neonatal intensive care unit, 1988-1997, and the impact of avoiding empiric vancomycin therapy. *Pediatrics* 2000;106:1387-90.
23. Stoll BJ, Hansen N, Fanaroff AA, Wright LL, Carlo WA, Ehrenkranz RA, et al. Late-onset sepsis in very low birth weight neonates: the experience of the NICHD Neonatal Research Network. *Pediatrics*. 2002;110:285-91.
24. Polin RA; Committee on Fetus and Newborn. Management of neonates with suspected or proven early-onset bacterial sepsis. *Pediatrics* 2012;129:1006-15.
25. May M, Daley AJ, Donath S, Isaacs D; Australasian Study Group for Neonatal Infections. Early onset neonatal meningitis in Australia and New Zealand, 1992-2002. *Arch Dis Child Fetal Neonatal Ed* 2005;90:F324-7.
26. Sgro M, Shah PS, Campbell D, Tenuta A, Shivananda S, Lee SK; Canadian Neonatal Network. Early-onset neonatal sepsis: rate and organism pattern between 2003 and 2008. *J Perinatol* 2011;31:794-8.
27. Engle WA, McGuire WA, Schreiner RL, Yu PL. Neutrophil storage pool depletion in neonates with sepsis and neutropenia. *J Pediatr* 1988;113:747-9.
28. Tolsma V, Schwebel C, Azoulay E, Darmon M, Souweine B, Vesin A, et al. Sepsis severe or septic shock: outcome according to immune status and immunodeficiency profile. *Chest* 2014;146:1205-1213.
29. Patel A, Gruber P. Severe infections in neutropenic patients. *Curr Opin Crit Care* 2015;21:586-92.

30. Goldstein B, Giroir B, Randolph A; International Consensus Conference on Pediatric Sepsis. International pediatric sepsis consensus conference: definitions for sepsis and organ dysfunction in pediatrics. *Pediatr Crit Care Med* 2005;6:2-8.
31. Wynn JL, Wong HR. Pathophysiology and treatment of septic shock in neonates. *Clin Perinatol* 2010;37:439-79.

Accepted Article



**Table 1.** Distribution of pathogens isolated from patients with early-onset sepsis (n=45)

Microorganisms	Term (n=14)	Preterm (n=31)	Total, n (%)
<b>Gram-positive bacteria</b>	<b>12</b>	<b>16</b>	<b>28 (62.2)</b>
<i>Streptococcus</i> group B	5	5	10 (22.2)
<i>Staphylococcus aureus</i>	4	1	5 (11.1)
<i>Staphylococcus epidermidis</i>	1	2	3 (6.7)
<i>Streptococcus mitis</i>	1	2	3 (6.7)
<i>Enterococcus faecalis</i>	1	1	2 (4.4)
<i>Enterococcus faecium</i>	0	2	2 (4.4)
<i>Staphylococcus capitis</i>	0	1	1 (2.2)
<i>Streptococcus anginosus</i>	0	1	1 (2.2)
<i>Streptococcus sanguinis</i>	0	1	1 (2.2)
<b>Gram-negative bacteria</b>	<b>2</b>	<b>14</b>	<b>16 (35.6)</b>
<i>Escherichia coli</i>	1	8	9 (20.0)
<i>Pseudomonas aeruginosa</i>	0	3	3 (6.7)
<i>Enterobacter aerogenes</i>	0	1	1 (2.2)
<i>Enterobacter cloacae</i>	0	1	1 (2.2)
<i>Haemophilus influenzae</i>	0	1	1 (2.2)
<i>Proteus mirabilis</i>	1	0	1 (2.2)
<b>Fungus</b>	<b>0</b>	<b>1</b>	<b>1 (2.2)</b>
<i>Candida albicans</i>	0	1	1 (2.2)

**Table 2.** Mortality and time of death in patients with early-onset sepsis

Patients with early-onset sepsis	Death	Time to death after infection		
		Days 1-3	Days 4-7	Days 8-14
n=45	17 (37.8)	8 (47.1)	4 (23.5)	5 (29.4)

Data are presented as n (%).

**Table 3.** Incidence of fatal infections by causative organisms (n=45)

Microorganisms	Case ratio	Percentage
<b>Gram-positive bacteria</b>	<b>6/28</b>	<b>21.4</b>
<i>Streptococci</i> other*	2/5	40.0
Coagulase-negative <i>staphylococci</i>	1/4	25.0
<i>Enterococci</i>	1/4	25.0
<i>Staphylococcus aureus</i>	1/5	20.0
<i>Streptococcus</i> group B	1/10	10.0
<b>Gram-negative bacteria</b>	<b>11/16</b>	<b>68.8</b>
<i>Pseudomonas aeruginosa</i>	3/3	100.0
<i>Haemophilus influenzae</i>	1/1	100.0
<i>Escherichia coli</i>	7/9	77.8
<i>Enterobacter</i> species	0/2	0.0
<i>Proteus mirabilis</i>	0/1	0.0
<b>Fungus</b>	<b>0/1</b>	<b>0.0</b>
<i>Candida albicans</i>	0/1	0.0

\*Defined as excluding *Streptococcus* group B

**Table 4.** Perinatal and clinical findings in patients with early-onset sepsis


Factors	Death (n=17)	Survival (n=28)	P value
<b>Perinatal characteristics</b>			
Gestational age (wk)	26.8±3.3	35.1±4.3	<0.001
Birth weight (g)	957.1±542.4	2,520.4±924.4	<0.001
Birth weight, <1,500 g	14 (82.4)	4 (14.3)	<0.001
Cesarean delivery	14 (82.4)	15 (53.6)	0.051
Sex (male /female)	11/6	15/13	0.463
Birth place (inborn/outborn)	17/0	22/6	0.069
1-min Apgar score	5.3±1.5	7.0±1.1	<0.001
5-min Apgar score	7.1±1.1	8.4±0.7	<0.001
Chorioamnionitis	15 (88.2)	8 (28.6)	<0.001
Antenatal steroid	10 (58.8)	3 (10.7)	0.001
Antenatal antibiotics	7 (41.2)	2 (7.1)	0.017
Maternal hypertension	2 (11.8)	2 (7.1)	0.626
Rupture of membranes, ≥18 hr	2 (11.8)	4 (14.3)	1.000
<b>Clinical Symptoms</b>			
Decreased activity	13 (76.5)	5 (17.9)	<0.001
Tachycardia, >160/min	13 (76.5)	16 (57.1)	0.189
Bradycardia, <100/min	8 (47.1)	4 (14.3)	0.034
Hypotension	10 (58.8)	0 (0)	<0.001
Apnea	3 (17.6)	4 (14.3)	1.000
Abdominal distension	2 (11.8)	3 (10.7)	1.000
Fever, ≥37.8 °C	1 (5.9)	10 (35.7)	0.033
<b>Laboratory Findings</b>			
WBC, ≥20,000/mm <sup>3</sup>	2 (11.8)	0 (0)	0.137
WBC, <5,000/mm <sup>3</sup>	8 (47.1)	7 (25)	0.128
Neutrophil, ≥7,000/mm <sup>3</sup>	3 (17.6)	8 (28.6)	0.493
Neutrophil, <1,000/mm <sup>3</sup>	7 (41.2)	1 (3.6)	0.003
Lymphocyte, ≥4,000/mm <sup>3</sup>	1 (5.9)	0 (0)	0.378
Lymphocyte, <1,000/mm <sup>3</sup>	4 (23.5)	9 (32.1)	0.737
Neutrophil-to-lymphocyte ratio	1.9±3.2	5.1±7.6	0.106
Platelet, <150,000/mm <sup>3</sup>	6 (35.3)	3 (10.7)	0.063
C-reactive protein, ≥0.5 mg/dL	9 (52.9)	20 (71.4)	0.209
<b>Comorbid diseases and empirical antibiotics</b>			
Respiratory distress syndrome	17 (100)	9 (32.1)	<0.001
Patent ductus arteriosus	6 (35.3)	0 (0)	0.002
Pulmonary hemorrhage	4 (23.5)	0 (0)	0.016
Intraventricular hemorrhage, ≥stage III	2 (11.8)	0 (0)	0.137
Acute kidney injury	1 (5.9)	0 (0)	0.378
Necrotizing enterocolitis, ≥stage II	1 (5.9)	0 (0)	0.378
Pneumothorax	1 (5.9)	0 (0)	0.378
Meningitis	0 (0)	2 (7.1)	0.519
Inadequate antimicrobial regimen	6 (35.3)	7 (25)	0.511

Data are presented as mean±SD or n (%).

**Table 5.** Logistic regression analysis adjusted for differences in gestational age and birth weight between the death and survival groups

Factors	$\beta$ -coefficient	OR (95% CI)	<i>P</i> value
Decreased activity	3.527	34.032 (1.829-633.369)	0.018
Gram-negative pathogens	3.745	42.289 (1.395-1281.829)	0.031
Neutropenia	3.830	46.057 (1.302-1628.646)	0.035
Antenatal steroid	-	-	0.178
Bradycardia	-	-	0.245
Fever	-	-	0.263
5-min Apgar score	-	-	0.557
Chorioamnionitis	-	-	0.716
1-min Apgar score	-	-	0.749
Antenatal antibiotics	-	-	0.840
Hypotension	-	-	0.998
Respiratory distress syndrome	-	-	0.998
Patent ductus arteriosus	-	-	0.999
Pulmonary hemorrhage	-	-	0.999

**Fig. 1.** Schematic diagram showing a 2-step classification for the study population. First, infants with identified microbes in their blood specimens were classified as infected or contaminated. Second, the infants with true infections were divided into the death and survival groups.



## Legends of Figures

**Fig. 1.** Schematic diagram showing a 2-step classification for the study population. First, infants with identified microbes in their blood specimens were classified as infected or contaminated. Second, the infants with true infections were divided into the death and survival groups.

Accepted Article